

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
30 November 2000 (30.11.2000)

PCT

(10) International Publication Number
WO 00/71514 A1

(51) International Patent Classification⁷: **C07C 317/48**,
317/50, 323/65, C07D 211/62, 213/56, 213/81, 213/82,
215/50, 295/15, 307/68, 317/58, C07K 5/062, A61K
31/10, A61P 7/00, 11/00

(21) International Application Number: **PCT/US00/09807**

(22) International Filing Date: **17 May 2000 (17.05.2000)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data: **60/134,996** **20 May 1999 (20.05.1999)** **US**

(71) Applicant (for all designated States except US): **G.D. SEARLE & CO.** [US/US]; Corporate Patent Department,
P.O. Box 5110, Chicago, IL 60680 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BARTA, Thomas**,
E. [US/US]; 1133 Maple Avenue, 3W, Evanston, IL 60202
(US). **BECKER, Daniel, P.** [US/US]; 1800 Maplewood
Lane, Glenview, IL 60025 (US). **BEDELL, Louis, J.**
[US/US]; 1832 E. Camp McDonald Road, Mt. Prospect,
IL 60056 (US). **DE CRESCENZO, Gary, A.** [US/US];
7345 Spruce Hill Court, St. Charles, MO 63304 (US).
FRESKOS, John, N. [US/US]; 7572 York, Clayton, MO
63105 (US). **GETMAN, Daniel, P.** [US/US]; 66 Sunny
Hill Court, Chesterfield, MO 63017 (US). **HEINTZ,**

Robert, M. [US/US]; 603 Nancy Place, Ballwin, MO
63021 (US). **HOCKERMAN, Susan, L.** [US/US];
5318 W. Hutchinson, Chicago, IL 60641 (US). **LI, Hui**
[US/US]; 322 Jefferson Court, Vernon Hills, MO 60061
(US). **MISCHKE, Brent, V.** [US/US]; 25 White River
Lane, Defiance, MO 63341 (US). **VILLAMIL, Clara, I.**
[US/US]; 813 Long Road, Glenview, IL 60025 (US).

(74) Agents: **WILLIAMS, Roger, A.** et al.; G.D. Searle & Co.,
Corporate Patent Department, P.O. Box 5110, Chicago, IL
60680-5110 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE,
DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU,
ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,
LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ,
PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT,
TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- With international search report.
- Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.

[Continued on next page]

(54) Title: **\$G(A)\$-AMINO-\$G(B)\$-SULFONYL HYDROXAMIC ACID COMPOUNDS**



(57) Abstract: A family of molecules of formula A in which R¹ is a 5- or 6-membered carbocyclo, heterocyclo, aryl or heteroaryl radical such that its size defined by certain parameters; R²⁰ is -O-R²¹, -NR¹³-O-R²² or -NR¹³-O-R¹⁴; and the other variables are as defined in the claims, is disclosed that inhibit matrix metalloproteinase (MMP) activity, and particularly inhibit the activity of one or more of MMP-2, MMP-9, or MMP-13, while generally exhibiting little activity against MMP-1. A contemplated compound also exhibits little inhibition of the production of TNF. Also disclosed are process for preparing a contemplated compound and for treating a mammal having a condition associated with pathological matrix metalloproteinase activity.

WO 00/71514 A1



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

α -AMINO- β -SULFONYL
HYDROXAMIC ACID COMPOUNDS

Description

5

Technical Field

This invention is directed to proteinase (protease) inhibitors, and more particularly to α -amino- β -sulfonyl carbocyclo, heterocyclo, aryl, or
10 heteroaryl hydroxamic acid compounds that, *inter alia*, inhibit the activity of matrix metalloproteinases, compositions of those inhibitors, intermediates for the syntheses of those compounds, processes for the preparation of the compounds and
15 processes for treating pathological conditions associated with pathological matrix metalloproteinase activity.

Background of the Invention

20 Connective tissue, extracellular matrix constituents and basement membranes are required components of all mammals. These components are the biological materials that provide rigidity, differentiation, attachments and, in some cases,
25 elasticity to biological systems including human beings and other mammals. Connective tissues components include, for example, collagen, elastin, proteoglycans, fibronectin and laminin. These biochemicals makeup, or are components of structures,
30 such as skin, bone, teeth, tendon, cartilage, basement membrane, blood vessels, cornea and vitreous humor.

Under normal conditions, connective tissue turnover and/or repair processes are controlled and

SUBSTITUTE SHEET (RULE 26)

in equilibrium. The loss of this balance for whatever reason leads to a number of disease states. Inhibition of the enzymes responsible loss of equilibrium provides a control mechanism for this tissue decomposition and, therefore, a treatment for these diseases.

Degradation of connective tissue or connective tissue components is carried out by the action of proteinase enzymes released from resident tissue cells and/or invading inflammatory or tumor cells. A major class of enzymes involved in this function is the zinc metalloproteinases (metalloproteases, or MMPs).

The metalloprotease enzymes are divided into classes with some members having several different names in common use. Examples are: collagenase I (MMP-1, fibroblast collagenase; EC 3.4.24.3); collagenase II (MMP-8, neutrophil collagenase; EC 3.4.24.34), collagenase III (MMP-13), stromelysin 1 (MMP-3; EC 3.4.24.17), stromelysin 2 (MMP-10; EC 3.4.24.22), proteoglycanase, matrilysin (MMP-7), gelatinase A (MMP-2, 72kDa gelatinase, basement membrane collagenase; EC 3.4.24.24), gelatinase B (MMP-9, 92kDa gelatinase; EC 3.4.24.35), stromelysin 3 (MMP-11), metalloelastase (MMP-12, HME, human macrophage elastase) and membrane MMP (MMP-14). MMP is an abbreviation or acronym representing the term Matrix Metalloprotease with the attached numerals providing differentiation between specific members of the MMP group.

The uncontrolled breakdown of connective tissue by metalloproteases is a feature of many pathological conditions. Examples include rheumatoid arthritis, osteoarthritis, septic arthritis; corneal,

epidermal or gastric ulceration; tumor metastasis, invasion or angiogenesis; periodontal disease; proteinuria; Alzheimer's Disease; coronary thrombosis and bone disease. Defective injury repair processes
5 can also occur. This can produce improper wound healing leading to weak repairs, adhesions and scarring. These latter defects can lead to disfigurement and/or permanent disabilities as with post-surgical adhesions.

10 Matrix metalloproteases are also involved in the biosynthesis of tumor necrosis factor (TNF) and inhibition of the production or action of TNF and related compounds is an important clinical disease treatment mechanism. TNF- α , for example, is a
15 cytokine that at present is thought to be produced initially as a 28 kD cell-associated molecule. It is released as an active, 17 kD form that can mediate a large number of deleterious effects *in vitro* and *in vivo*. For example, TNF can cause and/or contribute
20 to the effects of inflammation, rheumatoid arthritis, autoimmune disease, multiple sclerosis, graft rejection, fibrotic disease, cancer, infectious diseases, malaria, mycobacterial infection, meningitis, fever, psoriasis,
25 cardiovascular/pulmonary effects such as post-ischemic reperfusion injury, congestive heart failure, hemorrhage, coagulation, hyperoxic alveolar injury, radiation damage and acute phase responses like those seen with infections and sepsis and during
30 shock such as septic shock and hemodynamic shock. Chronic release of active TNF can cause cachexia and anorexia. TNF can be lethal.

TNF- α convertase is a metalloproteinase involved in the formation of active TNF- α .

SUBSTITUTE SHEET (RULE 26)

Inhibition of TNF- α convertase inhibits production of active TNF- α . Compounds that inhibit both MMPs activity have been disclosed in WIPO International Publication Nos. WO 94/24140, WO 94/02466, WO 5 97/20824 and EP 0 780 386 A1. There remains a need for effective MMP and TNF- α convertase inhibiting agents. Compounds that inhibit MMPs such as collagenase, stromelysin and gelatinase have been shown to inhibit the release of TNF (Gearing et al. 10 Nature 376, 555-557 (1994), McGeehan et al., Nature 376, 558-561 (1994)).

MMPs are involved in other biochemical processes in mammals as well. Included is the control of ovulation, post-partum uterine involution, 15 possibly implantation, cleavage of APP (β -Amyloid Precursor Protein) to the amyloid plaque and inactivation of α_1 -protease inhibitor (α_1 -PI).

Inhibition of these metalloproteases permits the control of fertility and the treatment or prevention 20 of Alzheimers Disease. In addition, increasing and maintaining the levels of an endogenous or administered serine protease inhibitor drug or biochemical such as α_1 -PI supports the treatment and prevention of diseases such as emphysema, pulmonary 25 diseases, inflammatory diseases and diseases of aging such as loss of skin or organ stretch and resiliency.

Inhibition of selected MMPs can also be desirable in other instances. Treatment of cancer and/or inhibition of metastasis and/or inhibition of 30 angiogenesis are examples of approaches to the treatment of diseases wherein the selective inhibition of stromelysin (MMP-3), gelatinase (MMP-2), gelatinase B (MMP-9) or collagenase III (MMP-13)

are the relatively most important enzyme or enzymes to inhibit especially when compared with collagenase I (MMP-1). A drug that does not inhibit collagenase I can have a superior therapeutic profile.

- 5 Osteoarthritis, another prevalent disease wherein it is believed that cartilage degradation in inflamed joints is at least partially caused by MMP-13 released from cells such as stimulated chondrocytes, may be best treated by administration of drugs one of
10 whose modes of action is inhibition of MMP-13. See, for example, Mitchell et al., *J. Clin. Invest.*, 97:761-763 (1996) and Reboul et al., *J. Clin. Invest.*, 97:2011-2019 (1996).

- Inhibitors of metalloproteases are known.
15 Examples include natural biochemicals such as tissue inhibitor of metalloproteinase (TIMP), α_2 -macroglobulin and their analogs or derivatives. These are high molecular weight protein molecules that form inactive complexes with metalloproteases.
20 A number of smaller peptide-like compounds that inhibit metalloproteases have been described. Mercaptoamide peptidyl derivatives have shown ACE inhibition *in vitro* and *in vivo*. Angiotensin converting enzyme (ACE) aids in the production of
25 angiotensin II, a potent pressor substance in mammals and inhibition of this enzyme leads to the lowering of blood pressure.

- Thiol group-containing amide or peptidyl amide-based metalloprotease (MMP) inhibitors are
30 known as is shown in, for example, WO95/12389, WO96/11209 and U.S. 4,595,700. Hydroxamate group-containing MMP inhibitors are disclosed in a number of published patent applications such as WO 95/29892, WO 97/24117, WO 97/49679 and EP 0 780 386 that

disclose carbon back-boned compounds, and WO 90/05719, WO 93/20047, WO 95/09841 and WO 96/06074 that disclose hydroxamates that have a peptidyl back-bones or peptidomimetic back-bones, as does the
5 article by Schwartz et al., *Progr. Med. Chem.*, 29:271-334 (1992) and those of Rasmussen et al., *Pharmacol. Ther.*, 75(1): 69-75 (1997) and Denis et al., *Invest. New Drugs*, 15(3): 175-185 (1997).

One possible problem associated with known
10 MMP inhibitors is that such compounds often exhibit the same or similar inhibitory effects against each of the MMP enzymes. For example, the peptidomimetic hydroxamate known as batimastat is reported to exhibit IC₅₀ values of about 1 to about 20 nanomolar
15 (nM) against each of MMP-1, MMP-2, MMP-3, MMP-7, and MMP-9. Marimastat, another peptidomimetic hydroxamate was reported to be another broad-spectrum MMP inhibitor with an enzyme inhibitory spectrum very similar to batimastat, except that marimastat
20 exhibited an IC₅₀ value against MMP-3 of 230 nM. Rasmussen et al., *Pharmacol. Ther.*, 75(1): 69-75 (1997).

Meta analysis of data from Phase I/II studies using marimastat in patients with advanced,
25 rapidly progressive, treatment-refractory solid tumor cancers (colorectal, pancreatic, ovarian, prostate) indicated a dose-related reduction in the rise of cancer-specific antigens used as surrogate markers for biological activity. Although marimastat
30 exhibited some measure of efficacy via these markers, toxic side effects were noted. The most common drug-related toxicity of marimastat in those clinical trials was musculoskeletal pain and stiffness, often commencing in the small joints in the hands,

spreading to the arms and shoulder. A short dosing holiday of 1-3 weeks followed by dosage reduction permits treatment to continue. Rasmussen et al., *Pharmacol. Ther.*, 75(1): 69-75 (1997). It is thought
5 that the lack of specificity of inhibitory effect among the MMPs may be the cause of that effect.

In view of the importance of hydroxamate MMP inhibitor compounds in the treatment of several diseases and the lack of enzyme specificity exhibited
10 by two of the more potent drugs now in clinical trials, it would be a great benefit if hydroxamates of greater enzyme specificity could be found. This would be particularly the case if the hydroxamate inhibitors exhibited strong inhibitory activity
15 against one or more of MMP-2, MMP-9 or MMP-13 that are associated with several pathological conditions, while at the same time exhibiting limited inhibition of MMP-1, an enzyme that is relatively ubiquitous and as yet not associated with any pathological
20 condition. It would also be beneficial if a compound that inhibited one or more of the above MMPs, while sparing MMP-1, also did not inhibit production of TNF. The disclosure that follows describes one family of hydroxamate MMP inhibitors that exhibit
25 those desirable activities


Brief Summary of the Invention

The present invention is directed to a family of molecules that among other properties
30 inhibit matrix metalloprotease (MMP) activity, and particularly inhibit the activity of one or more of MMP-2, MMP-9, or MMP-13, while generally exhibiting little activity against MMP-1. A contemplated compound also exhibits little inhibition of the

- 8 -

production of TNF. The present invention is also directed to processes for preparing a contemplated compound and for treating a mammal having a condition associated with pathological matrix metalloprotease activity.

Briefly, one embodiment of the present invention is directed to an α -amino- β -sulfonyl carbocyclo, heterocyclo, aryl, or heteroaryl hydroxamic acid compound that is active as a matrix metalloprotease enzyme inhibitor. That compound corresponds in structure to Formula A.



A

wherein

15

R^2 is selected from the group consisting of a hydrido, C_1 - C_4 hydrocarbyl, hydroxy- C_1 - C_4 hydrocarbyl, C_1 - C_4 hydrocarbyloxy, halo- C_1 - C_4 hydrocarbyl, C_1 - C_4 hydrocarbyloxymethyl, aryl, aryl C_1 - C_4 hydrocarbyl, aminomethyl, (N- C_1 - C_3 hydrocarbyl)aminomethyl, (N,N-di- C_1 - C_3 hydrocarbyl)aminomethyl, (N-morpholino)methyl, (N-pyrrolidino)methyl, or (N-thiomorpholino)methyl group. R^2 is preferably a hydrido, hydroxy, hydroxymethyl, methoxymethyl or methyl-N-morpholinyl group.

25

R^1 is a substituent that contains an optionally substituted 5- or 6-membered carbocyclo,

SUBSTITUTE SHEET (RULE 26)

heterocyclo, aryl or heteroaryl radical bonded directly to the depicted SO₂-group that has a length of an extended pentyl group and less than about that of an extended eicosyl group, and preferably having a length that is longer than an extended hexyl group and less than that of an extended stearyl group. In addition, R¹ defines a three-dimensional volume upon rotation about an axis drawn through the SO₂-bonded 1-position and the 4-position of a 6-membered ring radical or drawn through the SO₂-bonded 1-position and the center of 3,4-bond of a 5-membered ring radical, whose widest dimension in a direction transverse to the axis of rotation is about that of one furanyl ring to about that of two phenyl rings.

R⁶ and R⁷ are independently selected from the group consisting of a hydrido, alkylcarbonyl, alkoxy carbonyl, arylcarbonyl, aralkylcarbonyl, carboxyalkyl, carboxyalkylcarbonyl, alkoxy carbonyl-aminoalkylcarbonyl, heterocyclocarbonyl, heterocycloalkylcarbonyl, aryloxy carbonylaminoalkylcarbonyl, aralkyloxy carbonylaminoalkylcarbonyl, perfluoroalkylcarbonyl, trifluoromethylalkylcarbonyl, alkylthioalkylcarbonyl, arylthioalkylcarbonyl, aralkylthioalkylcarbonyl, heteroaralkylthioalkylcarbonyl, or a sulfoxide or sulfone of any of said thio substituents, alkylsulfonyl, arylsulfonyl, and an aminocarbonylalkyl or aminocarbonylalkyl-carbonyl group

wherein the aminocarbonylalkyl or aminocarbonylalkylcarbonyl nitrogen is (i) unsubstituted, or (ii) is the reacted amine of an amino acid, or (iii) substituted with one or two radicals selected from the group consisting of an alkyl, hydroxyalkyl, hydroxyheteroaralkyl,

SUBSTITUTE SHEET (RULE 26)

cycloalkyl, aralkyl, trifluoromethylalkyl, heterocycloalkyl, benzofused heterocycloalkyl, benzofused heterocycloalkyl, benzofused cycloalkyl, and an N,N-dialkylsubstituted-alkylaminoalkyl group, or (iv) that carboxamido nitrogen and two substituents bonded thereto together form a 5- to 8-membered heterocyclo, heteroaryl or benzofused heterocycloalkyl ring that is itself unsubstituted or substituted with one or two radicals independently selected from the group consisting of an alkyl, alkoxy carbonyl, nitro, heterocycloalkyl, hydroxy, hydroxycarbonyl, aryl, aralkyl, heteroaralkyl and an amino group, or

R⁶ and R⁷ together with the depicted nitrogen atom form a saturated or unsaturated cyclic imide substituent in which the ring contains five to eight atoms.

R²⁰ is (a) -O-R²¹, where R²¹ is selected from the group consisting of a hydrido, C₁-C₆-alkyl, aryl, ar-C₁-C₆-alkyl group and a pharmaceutically acceptable cation, (b) -NR¹³-O-R²² wherein R²² is a selectively removable protecting group such as a 2-tetrahydropyranyl, benzyl, p-methoxybenzyl (MOZ), carbonyl-C₁-C₆-alkoxy, trisubstituted silyl group or o-nitrophenyl group, peptide synthesis resin and the like, wherein trisubstituted silyl group is substituted with C₁-C₆-alkyl, aryl, or ar-C₁-C₆-alkyl, and R¹³ is a hydrido, C₁-C₆-alkyl or benzyl group, or (c) -NR¹³-O-R¹⁴, where R¹³ is as before and R¹⁴ is a hydrido, a pharmaceutically acceptable cation or C(W)R¹⁵ group, where W is O or S and R¹⁵ is

selected from the group consisting of an C₁-C₆-alkyl, aryl, C₁-C₆-alkoxy, heteroaryl-C₁-C₆-alkyl, C₃-C₈-cycloalkyl-C₁-C₆-alkyl, aryloxy, ar-C₁-C₆-alkoxy, ar-C₁-C₆-alkyl, heteroaryl and amino C₁-C₆-alkyl group
5 wherein the aminoalkyl nitrogen is (i) unsubstituted or (ii) substituted with one or two substituents independently selected from the group consisting of an C₁-C₆-alkyl, aryl, ar-C₁-C₆-alkyl, C₃-C₈-cycloalkyl-C₁-C₆-alkyl, ar-C₁-C₆-alkoxycarbonyl, C₁-
10 C₆-alkoxycarbonyl, and C₁-C₆-alkanoyl radical, or (iii) wherein the amino C₁-C₆-alkyl nitrogen and two substituents attached thereto form a 5- to 8-membered heterocyclo or heteroaryl ring.

A 5- or 6-membered carbocyclo, heterocyclo, aryl or heteroaryl radical bonded directly to the depicted SO₂-group that constitutes an R¹ group whose 5- or 6-membered carbocyclo, heterocyclo, aryl or heteroaryl radical of R¹ is itself optionally substituted with an -A-R-E-Y substituent. In such an
20 -A-R-E-Y substituent, A is selected from the group consisting of

- (1) -O-;
- (2) -S-;
- (3) -NR^k-;
- 25 (4) -CO-N(R^k) or -N(R^k)-CO-;
- (5) -CO-O- or -O-CO-;
- (6) -O-CO-O-;
- (7) -HC=CH-;
- (8) -NH-CO-NH-;
- 30 (9) -C≡C-;
- (10) -N=N-;
- (11) -NH-NH-;

SUBSTITUTE SHEET (RULE 26)

-12-

(12) $-\text{CS}-\text{N}(\text{R}^k)-$ or $-\text{N}(\text{R}^k)-\text{CS}-$;

(13) $-\text{CH}_2-$;

(14) $-\text{O}-\text{CH}_2-$ or $-\text{CH}_2-\text{O}-$;

(15) $-\text{S}-\text{CH}_2-$ or $-\text{CH}_2-\text{S}-$;

5 (16) $-\text{SO}-$; and

(17) $-\text{SO}_2-$; or

(18) A is absent and R is directly bonded to the depicted SO_2 -bonded 5- or 6-membered ring group.

10 The moiety R is selected from the group consisting of alkyl, alkoxy, halo, alkoxyalkyl, aryl, heteroaryl, cycloalkyl, heterocycloalkyl, aralkyl, heteroaralkyl, heterocycloalkyl, cycloalkyl, cycloalkoxyalkyl, heterocycloalkoxyalkyl, 15 aryloxyalkyl, heteroaryloxy-alkyl, arylthioalkyl, heteroarylthioalkyl, cycloalkylthioalkyl, and a heterocycloalkylthioalkyl group wherein the aryl, heteroaryl, cycloalkyl or heterocycloalkyl substituent is (i) unsubstituted or (ii) substituted 20 with one or two radicals selected from the group consisting of a halo, alkyl, perfluoroalkyl, perfluoroalkoxy, perfluoroalkylthio, trifluoromethylalkyl, amino, alkoxycarbonylalkyl, alkoxy, C_1 - C_2 -alkylenedioxy, hydroxycarbonylalkyl, 25 hydroxycarbonylalkylamino, nitro, hydroxy, hydroxyalkyl, alkanoylamino, and a alkoxycarbonyl group.

The group E is selected from the group consisting of

30 (1) $-\text{CO}(\text{R}^g)-$ or $-(\text{R}^g)\text{CO}-$;

(2) $-\text{CON}(\text{R}^k)-$ or $-(\text{R}^k)\text{NCO}-$;

(3) $-\text{CO}-$;

SUBSTITUTE SHEET (RULE 26)

-13-

(4) $-\text{SO}_2-\text{R}^g$ or $-\text{R}^g\text{SO}_2-$;(5) $-\text{SO}_2-$;(6) $-\text{N}(\text{R}^k)-\text{SO}_2-$ or $-\text{SO}_2-\text{N}(\text{R}^k)-$; or

(7) E is absent and R is bonded directly to Y; and

5

Y is absent or is selected from the group consisting of a hydrido, alkyl, alkoxy, haloalkyl, aryl, aralkyl, cycloalkyl, heteroaryl, hydroxy, nitrile, nitro, aryloxy, aralkoxy, heteroaryloxy,

10

heteroaralkyl, $\text{R}^c\text{oxyalkyl}$, perfluoroalkoxy, perfluoroalkylthio, trifluoromethylalkyl, alkenyl, heterocycloalkyl, cycloalkyl, trifluoromethyl, alkoxycarbonyl, and a aminoalkyl group, wherein the aryl, heteroaryl or heterocycloalkyl group is (i)

15

unsubstituted or (ii) substituted with one or two radicals independently selected from the group

consisting of an alkanoyl, halo, nitro, nitrile, haloalkyl, alkyl, aralkyl, aryl, alkoxy, and an amino group wherein the amino nitrogen is (i) unsubstituted

20

or (ii) substituted with one or two groups independently selected from hydrido, alkyl, and an aralkyl group.

Preferably, an $-\text{R}^1$ substituent is a substituent $-\text{G}-\text{A}-\text{R}-\text{E}-\text{Y}$ in which

25

G is an aryl or heteroaryl group;

the substituent A is selected from the group consisting of

(1) $-\text{O}-$;(2) $-\text{S}-$;

30

(3) $-\text{NR}^k-$;(4) $-\text{CO}-\text{N}(\text{R}^k)$ or $-\text{N}(\text{R}^k)-\text{CO}-$;(5) $-\text{CO}-\text{O}-$ or $-\text{O}-\text{CO}-$;(6) $-\text{O}-\text{CO}-\text{O}-$;**SUBSTITUTE SHEET (RULE 26)**

-14-

- (7) -HC=CH-;
- (8) -NH-CO-NH-;
- (9) -C≡C-;
- (10) -N=N-;
- 5 (11) -NH-NH-;
- (12) -CS-N(R^k)- or -N(R^k)-CS-;
- (13) -CH₂-;
- (14) -O-CH₂- or -CH₂-O-;
- (15) -S-CH₂- or -CH₂-S-;
- 10 (16) -SO-; and
- (17) -SO₂-; or
- (18) A is absent and R is directly bonded to the N-heterocyclo group, G.

The moiety R is selected from the group

15 consisting of alkyl, alkoxy, halo, alkoxyalkyl, aryl, heteroaryl, cycloalkyl, heterocycloalkyl, aralkyl, heteroaralkyl, heterocycloalkyl, cycloalkyl, cycloalkoxyalkyl, heterocycloalkoxyalkyl, aryloxyalkyl, heteroaryloxyalkyl, arylthioalkyl,

20 heteroarylthioalkyl, cycloalkylthioalkyl, and a heterocycloalkylthioalkyl group wherein the aryl or heteroaryl or cycloalkyl or heterocycloalkyl substituent is (i) unsubstituted or (ii) substituted with one or two radicals selected from the group

25 consisting of a halo, alkyl, perfluoroalkyl, perfluoroalkoxy, perfluoroalkylthio, trifluoromethylalkyl, amino, alkoxycarbonylalkyl, alkoxy, C₁-C₂-alkylenedioxy, hydroxycarbonylalkyl, hydroxycarbonylalkylamino, nitro, hydroxy,

30 hydroxyalkyl, alkanoylamino, and a alkoxycarbonyl group;

The moiety E is selected from the group consisting of

SUBSTITUTE SHEET (RULE 26)

-15-

- (1) $-\text{CO}(\text{R}^g)-$ or $-(\text{R}^g)\text{CO}-$;
- (2) $-\text{CON}(\text{R}^k)-$ or $-(\text{R}^k)\text{NCO}-$;
- (3) $-\text{CO}-$;
- (4) $-\text{SO}_2-\text{R}^g-$ or $-\text{R}^g-\text{SO}_2-$;
- 5 (5) $-\text{SO}_2-$;
- (6) $-\text{N}(\text{R}^k)-\text{SO}_2-$ or $-\text{SO}_2-\text{N}(\text{R}^k)-$; or
- (7) E is absent and R is bonded directly to Y; and
the moiety Y is absent or is selected from
the group consisting of a hydrido, alkyl, alkoxy,
10 haloalkyl, aryl, aralkyl, cycloalkyl, heteroaryl,
hydroxy, nitrile, nitro, aryloxy, aralkoxy,
heteroaryloxy, heteroaralkyl, R^{Coxy} alkyl,
perfluoroalkoxy, perfluoroalkylthio,
trifluoromethylalkyl, alkenyl, heterocycloalkyl,
15 cycloalkyl, trifluoromethyl, alkoxycarbonyl, and a
aminoalkyl group, wherein the aryl, heteroaryl or
heterocycloalkyl group is (i) unsubstituted or (ii)
substituted with one or two radicals independently
selected from the group consisting of an alkanoyl,
20 halo, nitro, nitrile, haloalkyl, alkyl, aralkyl,
aryl, alkoxy, and an amino group wherein the amino
nitrogen is (i) unsubstituted or (ii) substituted
with one or two groups independently selected from
hydrido, alkyl, and an aralkyl group; and wherein
25 R^d and R^e are independently selected from
the group consisting of a hydrido, alkyl, alkenyl,
alkenyl, arylalkyl, aryl, alkanoyl, aroyl,
arylalkylcarbonyl, alkoxycarbonyl or
arylalkyloxycarbonyl group;
- 30 R^f is selected from the group consisting of
a nitro, hydroxy, alkyl, halogen (halo; F, Cl, Br,
I), aryl, alkoxy, cyano, and R^dR^e amino group;

SUBSTITUTE SHEET (RULE 26)

-16-

R^g is selected from the group consisting of a hydrido, aryl, heteroaryl, heterocyclo, aroyl, alkanoyl, heteroaroyl, halogen (F, Cl, Br, I), cyano, aldehydo (CHO, formyl), hydroxy, amino, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, alkoxy, aryloxy, heteroaryloxy, alkenyloxy, alkynyloxy, alkoxyaryl, alkoxyheteroaryl, R^hRⁱ-amino, alkoxyalkyl, alkylenedioxy, aryloxyalkyl, perfluoroalkyl, trifluoroalkyl, alkylthio, arylthio, alkyloxycarbonyl, alkyloxycarbonyloxy, aryloxycarbonyl, arylalkyloxycarbonyl, arylalkyloxycarbonylamino, aryloxycarbonyloxy, carboxy, R^hRⁱ-aminocarbonyloxy, R^hRⁱ-aminocarbonyl, R^hRⁱ-aminoalkanoyl, hydroxyaminocarbonyl, R^hRⁱ-aminosulfonyl, R^hRⁱ-aminocarbonyl(R^h)amino, trifluoromethylsulfonyl-(R^h)amino, heteroarylsulfonyl(R^h)amino, arylsulfonyl(R^h)amino, arylsulfonyl(R^h)aminocarbonyl, alkylsulfonyl(R^h)amino, arylcarbonyl(R^h)aminosulfonyl, and an alkylsulfonyl(R^h)aminocarbonyl substituent;

wherein R^h is selected from the group consisting of an arylalkyl, aryl, heteroaryl, heterocyclo, alkyl, alkynyl, alkenyl, alkoxyalkyl, substituted or unsubstituted aminoalkyl, alkyloxycarbonyl, arylalkyloxycarbonyl, carboxyalkyl, haloalkyl, alkanoyl, aroyl, substituted or unsubstituted aminoalkanoyl, halo alkanoyl and a hydroxyalkyl group, each of which groups is optionally substituted by one or two groups independently selected from R^j substituents as are

SUBSTITUTE SHEET (RULE 26)

-17-

the substituents of the substituted aminoalkyl and substituted aminoalkanoyl groups;

R^i is selected from the group consisting of an arylalkyl, aryl, heteroaryl, heterocyclo, alkyl, alkynyl, alkenyl, alkoxyalkyl, alkoxyalkyl, substituted or unsubstituted aminoalkyl, alkyloxycarbonyl, arylalkyloxycarbonyl, carboxyalkyl, haloalkyl, alkanoyl, aroyl, substituted or unsubstituted aminoalkanoyl, halo alkanoyl and a hydroxyalkyl group, each of which groups are optionally substituted by one or two R^j substituents;


R^j is selected from the group consisting of an arylalkyl, aryl, heteroaryl, heterocyclo, alkyl, alkynyl, alkenyl, alkoxyalkyl, alkoxyalkyl, substituted or unsubstituted aminoalkyl, alkyloxycarbonyl, arylalkyloxycarbonyl, carboxyalkyl, haloalkyl, alkanoyl, aroyl, substituted or unsubstituted aminoalkanoyl, halo alkanoyl and a hydroxyalkyl group, wherein the substituents of the substituted aminoalkyl and substituted aminoalkanoyl groups are selected from the group consisting of an alkyl, alkenyl, alkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, aryloxycarbonyl and an alkyloxycarbonyl group; and

R^k is selected from hydrido, alkyl, alkenyl, alkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, aryloxycarbonyl, alkyloxycarbonyl, R^{CR^d} amino carbonyl, R^{CR^d} aminosulfonyl, R^{CR^d} aminoalkanoyl and R^{CR^d} aminoalkylsulfonyl, and R^a , R^c , R^d , and R^g are as defined hereinbefore.

A process for treating a host mammal having a condition associated with pathological matrix metalloprotease activity is also contemplated. That

SUBSTITUTE SHEET (RULE 26)

process comprises administering a compound corresponding in structure to Formula B that selectively inhibits one or more MMPs, while exhibiting less activity against at least MMP-1 in an
 5 MMP enzyme-inhibiting effective amount to a mammalian host having such a condition. A contemplated compound also does not substantially inhibit the production of TNF.



B


10

wherein R^1 , R^2 , R^6 , R^7 , R^{13} , and R^{14} are as defined before. Preferably, R^2 , R^6 , and R^{13} are hydrido, and R^7 is R^{7A} , wherein R^{7A} is selected from the group consisting of an alkylcarbonyl,
 15 alkoxycarbonyl, arylcarbonyl, alkylsulfonyl, arylsulfonyl, aminoalkylcarbonyl, aminoaralkylcarbonyl, alkoxycarbonylaminoalkylcarbonyl, alkoxycarbonylaminoaralkylcarbonyl, cycloalkoxycarbonylaminoalkylcarbonyl;
 20 cycloalkoxycarbonylaminoaralkylcarbonyl, aryloxycarbonylaminoalkylcarbonyl, aryloxycarbonylaminoaralkylcarbonyl, aralkyloxycarbonylaminoalkylcarbonyl, aralkyloxycarbonylaminoaralkylcarbonyl, heteroaryloxycarbonylaminoalkylcarbonyl and a
 25 heteroaryloxycarbonylaminoaralkylcarbonyl group.

An intermediate compound in the preparation of a particularly preferred compound of Formula A is also contemplated. Such a compound has a structure that corresponds to formula C, below.

SUBSTITUTE SHEET (RULE 26)


-19-



C

wherein R^2 , R^6 , R^7 and R^{20} are as before
 5 described and D is a nucleophilically-displacable
 leaving group such as $-F$ or $-NO_2$.

A particularly preferred intermediate for
 use in preparing a contemplated MMP inhibitor
 corresponds in structure to Formula D, below, wherein
 10 R^1 , R^2 , R^6 , R^7 and R^{13} are as before described.



D

Among the several benefits and advantages
 15 of the present invention are the provision of
 compounds and compositions that are effective for the
 inhibition of metalloproteinases implicated in
 diseases and disorders involving uncontrolled
 breakdown of connective tissue.

20 More particularly, a benefit of this
 invention is the provision of a compound and
 composition effective for inhibiting
 metalloproteinases, particularly MMP-13 and/or MMP-2,
 associated with pathological conditions such as, for

SUBSTITUTE SHEET (RULE 26)

example, rheumatoid arthritis, osteoarthritis, septic arthritis, corneal, epidermal or gastric ulceration, tumor metastasis, invasion or angiogenesis, periodontal disease, proteinuria, Alzheimer's
5 Disease, coronary thrombosis, plaque formation and bone disease.

An advantage of the invention is the provision of a method for preparing such compounds and compositions. Another benefit is the provision
10 of a method for treating a pathological condition associated with abnormal matrix metalloproteinase activity.

Another benefit of the invention is that a contemplated compound typically exhibits about equal
15 to greater inhibition of MMP-2 as compared to MMP-13, while maintaining sparing (lessened) inhibitory activity against MMP-1.

Another advantage of the invention is the provision of compounds, compositions and methods
20 effective for treating such pathological conditions by selective inhibition of a metalloproteinase such as MMP-13 and MMP-2 associated with such conditions with minimal side effects resulting from inhibition of other proteinases such as MMP-1, whose activity is
25 necessary or desirable for normal body function.

Yet another advantage of a contemplated compound is the selectivity with respect to tumor necrosis factor release and/or tumor necrosis factor receptor release that provides the physician with
30 another factor to help select the best drug for a particular patient. While not wishing to be bound by theory, it is believed that there are several factors to this type of selectivity to be considered.

The first is that presence of tumor necrosis factor can be desirable for the control of cancer in the organism, so long as TNF is not present in a toxic excess. Thus, uncontrolled inhibition of
5 release of TNF can be counterproductive and actually can be considered an adverse side effect even in cancer patients. In addition, selectivity with respect to inhibition of the release of the tumor necrosis factor receptor can also be desirable. The
10 presence of that receptor can be desirable for maintaining a controlled tumor necrosis level in the mammal by binding excess TNF.

Still further benefits and advantages of the invention will be apparent to the skilled worker
15 from the disclosure that follows.

Detailed Description of the Invention


In accordance with the present invention, it has been found that certain α -amino- β -sulfonyl
20 carbocyclo, heterocyclo, aryl or heteroaryl hydroxamic acids (hydroxamates) are effective, *inter alia*, for inhibition of matrix metalloproteinases ("MMPs") believed to be associated with uncontrolled or otherwise pathological breakdown of connective
25 tissue. In particular, it has been found that these certain α -amino- β -sulfonyl carbocyclo, heterocyclo, aryl or heteroaryl hydroxamic acid compounds are effective for inhibition of collagenase III (MMP-13) and also gelatinase A (MMP-2), which can be
30 particularly destructive to tissue if present or generated in abnormal quantities or concentrations, and thus exhibit a pathological activity.

Moreover, it has been discovered that many of these α -amino- β -sulfonyl carbocyclo, heterocyclo,

-22-

aryl or heteroaryl hydroxamic acids are selective in the inhibition of MMPs associated with diseased conditions without excessive inhibition of other collagenases essential to normal bodily function such as tissue turnover and repair. More particularly, it has been found that particularly preferred the α -amino- β -sulfonyl carbocyclo, heterocyclo, aryl or heteroaryl hydroxamic acid compounds are particularly active in inhibiting of MMP-13 and/or MMP-2, while having a limited or minimal effect on MMP-1. This point is discussed in detail hereinafter and is illustrated in the Inhibition Table hereinafter.

One embodiment of the present invention is directed to a α -amino- β -sulfonyl carbocyclo, heterocyclo, aryl or heteroaryl hydroxamic acid compound, a pharmaceutically acceptable salt of such a compound that can act as a matrix metalloprotease enzyme inhibitor, a precursor to such a compound or a pro-drug form of such a compound. A contemplated compound corresponds in structure to Formula A.



A

wherein

R² is selected from the group consisting of a hydrido, C₁-C₄ hydrocarbyl, hydroxy-C₁-C₄

SUBSTITUTE SHEET (RULE 26)

hydrocarbyl, C₁-C₄ hydrocarbyloxy, halo-C₁-C₄
hydrocarbyl, C₁-C₄ hydrocarbyloxymethyl, aryl, aryl
C₁-C₄ hydrocarbyl, aminomethyl, (N-C₁-C₃
hydrocarbyl)aminomethyl, (N,N-di-C₁-C₃
5 hydrocarbyl)aminomethyl, (N-morpholino)methyl, (N-
pyrrolidino)methyl, or (N-thiomorpholino)methyl
group. R² is preferably a hydrido, hydroxy,
hydroxymethyl, methoxymethyl, C₁-C₄ hydrocarbyl (e.g.
methyl), aryl (phenyl) or methyl-N-morpholinyl group,
10 with phenyl or C₁-C₄ hydrocarbyl being more
preferred;

R¹ is a substituent that contains an
optionally substituted 5- or 6-membered carbocyclo,
heterocyclo, aryl or heteroaryl radical bonded
15 directly to the depicted SO₂-group that has a length
of an extended pentyl group and less than about that
of an extended eicosyl group. In addition, R¹
defines a three-dimensional volume that upon rotation
about an axis drawn through the SO₂-bonded 1-position
20 and the 4-position of a 6-membered ring radical or
drawn through the SO₂-bonded 1-position and the
center of 3,4-bond of a 5-membered ring radical whose
widest dimension in a direction transverse to the
axis of rotation is about that of one furanyl ring to
25 about that of two phenyl rings; R¹ preferably has a
length that is greater than that of an extended
hexyl group and less than that of an extended stearyl
group.

R⁶ and R⁷ are independently selected from
30 the group consisting of a hydrido, alkylcarbonyl,
alkoxycarbonyl, arylcarbonyl, aralkylcarbonyl,
carboxyalkyl, carboxyalkylcarbonyl, alkoxycarbonyl-

aminoalkylcarbonyl, heterocyclocarbonyl,
heterocycloalkylcarbonyl, aryloxy-carbonylamino-
alkylcarbonyl, aralkyloxy-carbonylaminoalkylcarbonyl,
perfluoroalkylcarbonyl, trifluoromethylalkylcarbonyl,
5 alkylthioalkylcarbonyl, arylthioalkylcarbonyl,
aralkylthioalkylcarbonyl, heteroaralkylthio-
alkylcarbonyl, or a sulfoxide or sulfone of any of
said thio substituents, alkylsulfonyl, arylsulfonyl,
and an aminocarbonylalkyl or aminocarbonylalkyl-
10 carbonyl group

wherein the aminocarbonylalkyl or
aminocarbonylalkylcarbonyl nitrogen is (i)
unsubstituted, or (ii) is the reacted amine of
an amino acid, or (iii) substituted with one or
15 two radicals selected from the group consisting
of an alkyl, hydroxyalkyl, hydroxyheteroaralkyl,
cycloalkyl, aralkyl, trifluoromethylalkyl,
heterocycloalkyl, benzofused heterocycloalkyl,
benzofused heterocycloalkyl, benzofused
20 cycloalkyl, and an N,N-dialkylsubstituted-
alkylaminoalkyl group, or (iv) that carboxamido
nitrogen and two substituents bonded thereto
together form a 5- to 8-membered heterocyclo,
heteroaryl or benzofused heterocycloalkyl ring
25 that is itself unsubstituted or substituted with
one or two radicals independently selected from
the group consisting of an alkyl,
alkoxycarbonyl, nitro, heterocycloalkyl,
hydroxy, hydroxycarbonyl, aryl, aralkyl,
30 heteroaralkyl and an amino group, or

R^6 and R^7 together with the depicted
nitrogen atom form a saturated or unsaturated cyclic
imide substituent in which the ring contains five to
eight atoms;

SUBSTITUTE SHEET (RULE 26)

R^6 is preferably hydrido, and R^7 is preferably an alkylcarbonyl, alkoxycarbonyl, arylcarbonyl, alkylsulfonyl, arylsulfonyl, aminoalkylcarbonyl or a substituted aminoalkylcarbonyl group such as an alkoxycarbonyl-aminoalkylcarbonyl, cycloalkoxycarbonylaminoalkylcarbonyl, aryloxycarbonylaminoalkylcarbonyl, aralkyloxycarbonylaminoalkylcarbonyl, or heteroaryloxycarbonylaminoalkylcarbonyl group;

10 R^{20} is (a) $-O-R^{21}$, where R^{21} is selected from the group consisting of a hydrido, C_1 - C_6 -alkyl, aryl, ar- C_1 - C_6 -alkyl group and a pharmaceutically acceptable cation, (b) $-NR^{13}-O-R^{22}$ wherein R^{22} is a selectively removable protecting group such as a 2-tetrahydropyranyl, benzyl, p-methoxybenzyl (MOZ), carbonyl- C_1 - C_6 -alkoxy, trisubstituted silyl group or o-nitrophenyl group, peptide synthesis resin and the like, wherein trisubstituted silyl group is substituted with C_1 - C_6 -alkyl, aryl, or ar- C_1 - C_6 -alkyl, and R^{13} is a hydrido, C_1 - C_6 -alkyl or benzyl group, or (c) $-NR^{13}-O-R^{14}$, where R^{13} is as before and R^{14} is a hydrido, a pharmaceutically acceptable cation or $C(W)R^{15}$ group, where W is O or S and R^{15} is selected from the group consisting of an C_1 - C_6 -alkyl, aryl, C_1 - C_6 -alkoxy, heteroaryl- C_1 - C_6 -alkyl, C_3 - C_8 -cycloalkyl- C_1 - C_6 -alkyl, aryloxy, ar- C_1 - C_6 -alkoxy, ar- C_1 - C_6 -alkyl, heteroaryl and amino C_1 - C_6 -alkyl group wherein the aminoalkyl nitrogen is (i) unsubstituted or (ii) substituted with one or two substituents independently selected from the group consisting of an C_1 - C_6 -alkyl, aryl, ar- C_1 - C_6 -alkyl, C_3 - C_8 -

15
20
25
30

cycloalkyl-C₁-C₆-alkyl, ar-C₁-C₆-alkoxycarbonyl, C₁-C₆-alkoxycarbonyl, and C₁-C₆-alkanoyl radical, or (iii) wherein the amino C₁-C₆-alkyl nitrogen and two substituents attached thereto form a 5- to 8-membered heterocyclo or heteroaryl ring;

A 5- or 6-membered carbocyclo, heterocyclo, aryl or heteroaryl radical bonded directly to the depicted SO₂-group that constitutes an R¹ group that is itself optionally substituted with an -A-R-E-Y substituent. In such an -A-R-E-Y substituent, A is selected from the group consisting of

- (1) -O-;
- (2) -S-;
- (3) -NR^k-;
- (4) -CO-N(R^k) or -N(R^k)-CO-;
- (5) -CO-O- or -O-CO-;
- (6) -O-CO-O-;
- (7) -HC=CH-;
- (8) -NH-CO-NH-;
- (9) -C≡C-;
- (10) -N=N-;
- (11) -NH-NH-;
- (12) -CS-N(R^k) - or -N(R^k)-CS-;
- (13) -CH₂-;
- (14) -O-CH₂- or -CH₂-O-;
- (15) -S-CH₂- or -CH₂-S-;
- (16) -SO-; and
- (17) -SO₂-; or
- (18) A is absent and R is directly bonded to the depicted SO₂-bonded 5- or 6-membered ring group;

The moiety R is selected from the group consisting of alkyl, alkoxy, halo, alkoxyalkyl, aryl, heteroaryl, cycloalkyl, heterocycloalkyl, aralkyl, heteroaralkyl, heterocycloalkyl, cycloalkylalkyl, cycloalkoxyalkyl, heterocycloalkoxyalkyl, aryloxyalkyl, heteroaryloxy-alkyl, arylthioalkyl, heteroarylthioalkyl, cycloalkylthioalkyl, and a heterocycloalkylthioalkyl group wherein the aryl, heteroaryl, cycloalkyl or heterocycloalkyl substituent is (i) unsubstituted or (ii) substituted with one or two radicals selected from the group consisting of a halo, alkyl, perfluoroalkyl, perfluoroalkoxy, perfluoroalkylthio, trifluoromethylalkyl, amino, alkoxycarbonylalkyl, alkoxy, C₁-C₂-alkylenedioxy, hydroxycarbonylalkyl, hydroxycarbonylalkylamino, nitro, hydroxy, hydroxyalkyl, alkanoylamino, and a alkoxycarbonyl group.

The group E is selected from the group consisting of

- (1) -CO(R^G) - or -(R^G)CO-;
- (2) -CON(R^k) - or -(R^k)NCO-;
- (3) -CO-;
- (4) -SO₂-R^G or -R^GSO₂-;
- (5) -SO₂-;
- (6) -N(R^k)-SO₂- or -SO₂-N(R^k)-; or
- (7) E is absent and R is bonded directly to Y; and

Y is absent or is selected from the group consisting of a hydrido, alkyl, alkoxy, haloalkyl, aryl, aralkyl, cycloalkyl, heteroaryl, hydroxy, nitrile, nitro, aryloxy, aralkoxy, heteroaryloxy, heteroaralkyl, R^Coxyalkyl, perfluoroalkoxy,

SUBSTITUTE SHEET (RULE 26)

perfluoroalkylthio, trifluoromethylalkyl, alkenyl,
heterocycloalkyl, cycloalkyl, trifluoromethyl,
alkoxycarbonyl, and a aminoalkyl group, wherein the
aryl, heteroaryl or heterocycloalkyl group is (i)
5 unsubstituted or (ii) substituted with one or two
radicals independently selected from the group
consisting of an alkanoyl, halo, nitro, nitrile,
haloalkyl, alkyl, aralkyl, aryl, alkoxy, and an amino
group wherein the amino nitrogen is (i) unsubstituted
10 or (ii) substituted with one or two groups
independently selected from hydrido, alkyl, and an
aralkyl group.

Preferably, an $-R^1$ substituent is a
substituent -G-A-R-E-Y in which
15 G is an aryl or heteroaryl group;
the substituent A is selected from the
group consisting of

- (1) -O-;
- (2) -S-;
- 20 (3) $-NR^k-$;
- (4) $-\text{CO}-N(R^k)$ or $-N(R^k)-\text{CO}-$;
- (5) $-\text{CO}-\text{O}-$ or $-\text{O}-\text{CO}-$;
- (6) $-\text{O}-\text{CO}-\text{O}-$;
- (7) $-\text{HC}=\text{CH}-$;
- 25 (8) $-\text{NH}-\text{CO}-\text{NH}-$;
- (9) $-\text{C}\equiv\text{C}-$;
- (10) $-\text{N}=\text{N}-$;
- (11) $-\text{NH}-\text{NH}-$;
- (12) $-\text{CS}-N(R^k)-$ or $-N(R^k)-\text{CS}-$;
- 30 (13) $-\text{CH}_2-$;
- (14) $-\text{O}-\text{CH}_2-$ or $-\text{CH}_2-\text{O}-$;
- (15) $-\text{S}-\text{CH}_2-$ or $-\text{CH}_2-\text{S}-$;
- (16) $-\text{SO}-$; and

-29-

(17) $-\text{SO}_2-$; or

(18) A is absent and R is directly bonded to the N-heterocyclo group, G;

The moiety R is selected from the group
5 consisting of alkyl, alkoxy, halo, alkoxyalkyl, aryl, heteroaryl, cycloalkyl, heterocycloalkyl, aralkyl, heteroaralkyl, heterocycloalkyl, cycloalkylalkyl, cycloalkoxyalkyl, heterocycloalkoxyalkyl, aryloxyalkyl, heteroaryloxyalkyl, arylthioalkyl,
10 heteroarylthioalkyl, cycloalkylthioalkyl, and a heterocycloalkylthioalkyl group wherein the aryl or heteroaryl or cycloalkyl or heterocycloalkyl substituent is (i) unsubstituted or (ii) substituted with one or two radicals selected from the group
15 consisting of a halo, alkyl, perfluoroalkyl, perfluoroalkoxy, perfluoroalkylthio, trifluoromethylalkyl, amino, alkoxycarbonylalkyl, alkoxy, C_1 - C_2 -alkylenedioxy, hydroxycarbonylalkyl, hydroxycarbonylalkylamino, nitro, hydroxy,
20 hydroxyalkyl, alkanoylamino, and a alkoxycarbonyl group;

The moiety E is selected from the group consisting of

- 25 (1) $-\text{CO}(\text{R}^g)-$ or $-(\text{R}^g)\text{CO}-$;
(2) $-\text{CON}(\text{R}^k)-$ or $-(\text{R}^k)\text{NCO}-$;
(3) $-\text{CO}-$;
(4) $-\text{SO}_2-\text{R}^g-$ or $-\text{R}^g-\text{SO}_2-$;
(5) $-\text{SO}_2-$;
(6) $-\text{N}(\text{R}^k)-\text{SO}_2-$ or $-\text{SO}_2-\text{N}(\text{R}^k)-$; or
30 (7) E is absent and R is bonded directly to Y; and
the moiety Y is absent or is selected from the group consisting of a hydrido, alkyl, alkoxy,

SUBSTITUTE SHEET (RULE 26)

haloalkyl, aryl, aralkyl, cycloalkyl, heteroaryl,
hydroxy, nitrile, nitro, aryloxy, aralkoxy,
heteroaryloxy, heteroaralkyl, R^coxyalkyl,
perfluoroalkoxy, perfluoroalkylthio,
5 trifluoromethylalkyl, alkenyl, heterocycloalkyl,
cycloalkyl, trifluoromethyl, alkoxycarbonyl, and a
aminoalkyl group, wherein the aryl, heteroaryl or
heterocycloalkyl group is (i) unsubstituted or (ii)
substituted with one or two radicals independently
10 selected from the group consisting of an alkanoyl,
halo, nitro, nitrile, haloalkyl, alkyl, aralkyl,
aryl, alkoxy, and an amino group wherein the amino
nitrogen is (i) unsubstituted or (ii) substituted
with one or two groups independently selected from
15 hydrido, alkyl, and an aralkyl group; and wherein
R^d and R^e are independently selected from
the group consisting of a hydrido, alkyl, alkenyl,
alkenyl, arylalkyl, aryl, alkanoyl, aroyl,
arylalkylcarbonyl, alkoxycarbonyl or
20 arylalkyloxycarbonyl group;
R^f is selected from the group consisting of
a nitro, hydroxy, alkyl, halogen (halo; F, Cl, Br,
I), aryl, alkoxy, cyano, and R^dR^eamino group;
R^g is selected from the group consisting of
25 a hydrido, aryl, heteroaryl, heterocyclo, aroyl,
alkanoyl, heteroaroyl, halogen (F, Cl, Br, I), cyano,
aldehydo (CHO, formyl), hydroxy, amino, alkyl,
alkenyl, alkynyl, cycloalkyl, cycloalkenyl, alkoxy,
aryloxy, heteroaryloxy, alkenyloxy, alkynyloxy,
30 alkoxyaryl, alkoxyheteroaryl, R^hRⁱ-amino,
alkoxyalkyl, alkylenedioxy, aryloxyalkyl,
perfluoroalkyl, trifluoroalkyl, alkylthio, arylthio,
alkyloxycarbonyl, alkyloxycarbonyloxy,

aryloxycarbonyl, arylalkyloxycarbonyl,
arylalkyloxycarbonylamino, aryloxycarbonyloxy,
carboxy, R^hR^i -aminocarbonyloxy, R^hR^i -aminocarbonyl,
 R^hR^i -aminoalkanoyl, hydroxyaminocarbonyl, R^hR^i -
5 aminosulfonyl, R^hR^i -aminocarbonyl (R^h) amino,
trifluoromethylsulfonyl- (R^h) amino,
heteroarylsulfonyl (R^h) amino, arylsulfonyl (R^h) amino,
arylsulfonyl (R^h) aminocarbonyl,
alkylsulfonyl (R^h) amino,
10 arylcarbonyl (R^h) aminosulfonyl, and an
alkylsulfonyl (R^h) aminocarbonyl substituent;

wherein R^h is selected from the group
consisting of an arylalkyl, aryl, heteroaryl,
heterocyclo, alkyl, alkynyl, alkenyl, alkoxyalkyl,
15 alkoxyalkyl, substituted or unsubstituted aminoalkyl,
alkyloxycarbonyl, arylalkyloxycarbonyl, carboxyalkyl,
haloalkyl, alkanoyl, aroyl, substituted or
unsubstituted aminoalkanoyl, halo alkanoyl and a
hydroxyalkyl group, each of which groups is
20 optionally substituted by one or two groups
independently selected from R^j substituents as are
the substituents of the substituted aminoalkyl and
substituted aminoalkanoyl groups;


R^i is selected from the group consisting of
25 an arylalkyl, aryl, heteroaryl, heterocyclo, alkyl,
alkynyl, alkenyl, alkoxyalkyl, alkoxyalkyl,
substituted or unsubstituted aminoalkyl,
alkyloxycarbonyl, arylalkyloxycarbonyl, carboxyalkyl,
haloalkyl, alkanoyl, aroyl, substituted or
30 unsubstituted aminoalkanoyl, halo alkanoyl and a
hydroxyalkyl group, each of which groups are
optionally substituted by one or two R^j substituents;

-32-

R^j is selected from the group consisting of an arylalkyl, aryl, heteroaryl, heterocyclo, alkyl, alkynyl, alkenyl, alkoxyalkyl, alkoxyalkyl, substituted or unsubstituted aminoalkyl, alkyloxycarbonyl, arylalkyloxycarbonyl, carboxyalkyl, haloalkyl, alkanoyl, aroyl, substituted or unsubstituted aminoalkanoyl, halo alkanoyl and a hydroxyalkyl group, wherein the substituents of the substituted aminoalkyl and substituted aminoalkanoyl groups are selected from the group consisting of an alkyl, alkenyl, alkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, aryloxycarbonyl and an alkyloxycarbonyl group; and

R^k is selected from hydrido, alkyl, alkenyl, alkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, aryloxycarbonyl, alkyloxycarbonyl, R^CR^damino carbonyl, R^CR^daminosulfonyl, R^CR^daminoalkanoyl and R^CR^daminoalkylsulfonyl, and R^a, R^c, R^d, and R^g are as defined hereinbefore.

A more preferred group of contemplated compounds corresponds in structure to Formula B1, below,



B1


25

wherein R¹ and R¹⁴ are as defined before, and R^{7A} is selected from the group consisting of an alkylcarbonyl, alkoxycarbonyl, arylcarbonyl,

SUBSTITUTE SHEET (RULE 26)

alkylsulfonyl, arylsulfonyl, aminoalkylcarbonyl,
 aminoaralkylcarbonyl, alkoxycarbonylamino-
 alkylcarbonyl, alkoxycarbonylaminoaralkylcarbonyl,
 cycloalkoxycarbonylaminoalkylcarbonyl,
 5 cycloalkoxycarbonylaminoaralkylcarbonyl,
 aryloxycarbonylaminoalkylcarbonyl, aryloxycarbonyl-
 aminoaralkylcarbonyl, aralkyloxycarbonylamino-
 alkylcarbonyl, aralkyloxycarbonylaminoalkylcarbonyl,
 heteroaryloxycarbonylaminoalkylcarbonyl and a
 10 heteroaryloxycarbonylaminoaralkylcarbonyl group.

An intermediate compound useful in the
 preparation of a particularly preferred compound of
 Formula A is also contemplated. One such
 intermediate has a structure that corresponds to
 15 formula C, below.




C

wherein R^2 , R^6 , R^7 and R^{20} are as before
 20 described and D is a nucleophilically-displacable
 leaving group such as -F or -NO₂. Here, R^{20} is
 usually -O- R^{21} , wherein R^{21} is typically hydrido.

Another particularly preferred intermediate
 for use in preparing a contemplated MMP inhibitor
 25 corresponds in structure to Formula D, below, wherein
 R^1 , R^2 , R^6 , R^7 and R^{13} are as before described.

- 34 -



Preferably in Formula D, R^2 , R^6 and R^{13} are preferably hydrido, and R^7 is preferably R^{7A} .

5 When examined along its longest chain of atoms, an R^1 substituent, including its own substituent when present, has a total length that is greater than that of a fully extended saturated chain of five carbon atoms (a pentyl group); i.e., a length
10 of a hexyl chain in staggered conformation or longer. An R^1 substituent also has a length that is less than that of a fully extended saturated chain of about 20 carbons (an eicosyl group). Preferably, that length is greater than that of a hexyl group and shorter
15 than that of a stearyl group, even though many more atoms may be present in ring structures or substituents.

Looked at more generally, and aside from specific moieties from which it is constructed, an R^1
20 substituent (radical, group or moiety) has a length of a pentyl group or greater. Such an R^1 substituent also has a length that is less than that of an eicosyl group. That is to say that a R^1 is a substituent having a length greater than that of a
25 fully extended saturated five carbon chain and shorter than that of a fully extended saturated twenty carbon chain, and more preferably, a length greater than that of a hexyl group and less than that

SUBSTITUTE SHEET (RULE 26)

of a stearyl group. The radical chain lengths are measured along the longest linear atom chain in the radical, following the skeletal atoms of a ring where necessary. Each atom in the chain, e.g. carbon,
5 oxygen or nitrogen, is presumed to be carbon for ease in calculation.

Such lengths can be readily determined by using published bond angles, bond lengths and atomic radii, as needed, to draw and measure a chain, or by
10 building models using commercially available kits whose bond angles, lengths and atomic radii are in accord with accepted, published values. Radical (substituent) lengths can also be determined somewhat less exactly by presuming, as is done here, that all
15 atoms have bond lengths of saturated carbon, that unsaturated and aromatic bonds have the same lengths as saturated bonds and that bond angles for unsaturated bonds are the same as those for saturated bonds, although the above-mentioned modes of
20 measurement are preferred. For example, a 4-phenyl or 4-pyridyl group has a length of a four carbon chain, as does a propoxy group, whereas a 4-halophenyl group has the length of a five carbon chain and biphenyl group has the length of about an
25 eight carbon chain using a contemplated measurement mode.

In addition, an R^1 substituent, upon rotation about an axis drawn through the SO_2 -bonded 1-position and the 4-position of a 6-membered ring
30 radical or the SO_2 -bonded 1-position and through the 3,4 bond of a 5-membered ring radical defines a three-dimensional volume whose widest dimension has the width of about one furanyl ring to about the

-36-

width of two phenyl rings in a direction transverse to that axis to rotation.

When utilizing this width or volume criterion, a fused ring system such as a naphthyl or purinyl radical is considered to be a 6- or 5-membered ring that is substituted at appropriate positions numbered from the SO₂-linkage that is deemed to be at the 1-position as discussed before. Thus, a 2-naphthyl substituent or an 8-purinyl

substituent is an appropriately sized R¹ radical as to width when examined using the above rotational width criterion. On the other hand, a 1-naphthyl group or a 7- or 9-purinyl group is too large upon rotation and is excluded.

As a consequence of these length and width requirements, R¹ substituents such as 4-(phenyl)phenyl [biphenyl], 4-(4'-methoxyphenyl)-phenyl, 4-(phenoxy)phenyl, 4-(thiophenyl)phenyl [4-(phenylthio)phenyl], 4-(phenylazo)phenyl 4-(phenylureido)phenyl, 4-(anilino)phenyl, 4-(nicotinamido)phenyl, 4-(isonicotinamido)phenyl, 4-(picolinamido)phenyl and 4-(benzamido)phenyl are among particularly preferred R¹ substituents, with 4-(phenoxy)phenyl and 4-(thiophenyl)phenyl being most preferred.

An SO₂-linked cyclohydrocarbyl, heterocyclo, aryl or heteroaryl radical is a 5- or 6-membered single-ring that is itself substituted with one other substituent, R⁴. The SO₂-linked single-ringed cyclohydrocarbyl, heterocyclo, aryl or heteroaryl radical is R⁴-substituted at its own 4-position when a 6-membered ring and at its own 3- or 4-position when a 5-membered ring. The

SUBSTITUTE SHEET (RULE 26)

cyclohydrocarbyl, heterocyclo, aryl or heteroaryl radical to which R^4 is bonded in some embodiments is preferably a phenyl group, so that R^1 is preferably PhR^4 in which R^4 is bonded at the 4-position of the
5 SO_2 -linked phenyl (Ph) radical, and in which R^4 can itself be optionally substituted as is discussed hereinafter. In other embodiments, a heterocyclo or heteroaryl radical is preferred over a phenyl radical, with the R^4 substituent being linked at the
10 4-position relative to the bond between the ring and the SO_2 group.


A contemplated R^4 substituent can be a single-ringed cyclohydrocarbyl, heterocyclo, aryl or heteroaryl group or another substituent having a
15 chain length of 3 to about 14 carbon atoms such as a hydrocarbyl or hydrocarbyloxy group [e.g., C_3 - C_{14} hydrocarbyl or O - C_2 - C_{14} hydrocarbyl], a phenyl group, a phenoxy group [$-OC_6H_5$], a thiophenoxy group [phenylsulfanyl; $-SC_6H_5$], an anilino group [$-NHC_6H_5$],
20 a phenylazo group [$-N_2C_6H_5$], a phenylureido group [aniline carbonylamino; $-NHC(O)NH-C_6H_5$], a benzamido group [$-NHC(O)C_6H_5$], a nicotinamido group [$3-NHC(O)C_5H_4N$], an isonicotinamido group [$4-NHC(O)C_5H_4N$], or a picolinamido group
25 [$2-NHC(O)C_5H_4N$]. Additionally contemplated R^4 substituent groups include a heterocyclo, heterocyclohydrocarbyl, arylhydrocarbyl, arylheterocyclohydrocarbyl, heteroarylhydrocarbyl, heteroarylheterocyclo-hydrocarbyl,
30 arylhydrocarbyloxyhydrocarbyl, aryloxyhydrocarbyl, hydrocarboylhydrocarbyl, arylhydrocarboylhydrocarbyl,

- arylcarbonylhydrocarbyl, arylazoaryl,
arylhydrazinoaryl, hydrocarbyl-thiohydrocarbyl,
hydrocarbylthioaryl, arylthiohydrocarbyl,
heteroarylthiohydrocarbyl,
5 hydrocarbylthioarylhydrocarbyl, arylhydrocarbyl-
thiohydrocarbyl, arylhydrocarbylthioaryl,
arylhydrocarbylamino, heteroarylhydrocarbylamino, or
a heteroarylthio group.

- A contemplated R⁴ substituent can itself
10 also be substituted with one or more substituent
radicals at the meta- or para-position or both of a
six-membered ring with a single atom or a substituent
containing a longest chain of up to ten atoms,
excluding hydrogen. Exemplary substituent radicals
15 include a halo, hydrocarbyl, hydrocarbyloxy, nitro,
cyano, perfluorohydrocarbyl, trifluoromethyl-
hydrocarbyl, hydroxy, mercapto, hydroxycarbonyl,
aryloxy, arylthio, arylamino, arylhydrocarbyl, aryl,
heteroaryloxy, heteroarylthio, heteroarylamino,
20 heteroarhydrocarbyl, hydrocarbyloxycarbonyl-
hydrocarbyl, heterocyclooxy, hydroxycarbonyl-
hydrocarbyl, heterocyclothio, heterocycloamino,
cyclohydrocarbyloxy, cyclohydrocarbylthio,
cyclohydrocarbylamino, heteroarylhydrocarbyloxy,
25 heteroarylhydrocarbylthio, heteroaryl-
hydrocarbylamino, arylhydrocarbyloxy,
arylhydrocarbylthio, arylhydrocarbylamino,
heterocyclic, heteroaryl, hydroxycarbonyl-
hydrocarbyloxy, alkoxycarbonylalkoxy, hydrocarbyloyl,
30 arylcarbonyl, arylhydrocarbyloyl, hydrocarbyloxy,
arylhydrocarbyloxy, hydroxyhydrocarbyl, hydroxy-
hydrocarbyloxy, hydrocarbylthio, hydrocarbyloxy-
hydrocarbylthio, hydrocarbyloxycarbonyl,
hydroxycarbonylhydrocarbyloxy, hydrocarbyloxy-

carbonylhydrocarbyl, hydrocarbylhydroxycarbonyl-
 hydrocarbylthio, hydrocarbyloxycarbonyl-
 hydrocarbyloxy, hydrocarbyloxycarbonyl-
 hydrocarbylthio, amino, hydrocarbylcarbonylamino,
 5 arylcarbonylamino, cyclohydrocarbylcarbonylamino,
 heterocyclohydrocarbylcarbonylamino,
 arylhydrocarbylcarbonylamino, heteroaryl-
 carbonylamino, heteroarylhydrocarbylcarbonylamino,
 heterocyclohydrocarbyloxy, hydrocarbylsulfonylamino,
 10 arylsulfonylamino, arylhydrocarbylsulfonylamino,
 heteroarylsulfonylamino, heteroarylhydrocarbyl-
 sulfonylamino, cyclohydrocarbylsulfonylamino,
 heterocyclohydrocarbylsulfonylamino and N-
 monosubstituted or N,N-disubstituted aminohydrocarbyl
 15 group wherein the substituent(s) on the nitrogen are
 selected from the group consisting of hydrocarbyl,
 aryl, arylhydrocarbyl, cyclohydrocarbyl,
 arylhydrocarbyloxycarbonyl, hydrocarbyloxycarbonyl,
 and hydrocarbonyl, or wherein the nitrogen and two
 20 substituents attached thereto form a 5- to 8-membered
 heterocyclic or heteroaryl ring group.

Compounds that are intermediates in the
 preparation of a MMP-inhibiting compound are also
 contemplated by this invention. One preferred
 25 intermediate corresponds in structure to Formula C,
 below,




C

-40-

wherein R^2 , R^6 , R^7 and R^{20} are as before described and D is a nucleophilically-displacable leaving group such as $-F$ or $-NO_2$. R^2 and R^6 are preferably hydrido here, R^7 is preferably R^{7A} , and
5 R^{20} is preferably $-O-R^{21}$, wherein R^{21} is hydrido.

Another preferred intermediate compound corresponds in structure to Formula D, below, structure to Formula D, below, wherein R^1 , R^2 , R^6 , R^7 and R^{13} are as before described.

10











D

In preferred practice for a compound of Formula D, R^2 , R^6 and R^{13} are hydrido, and R^7 is R^{7A} .

15 Table 1 through Table 9, below, illustrate several synthesized compounds that are contemplated by this invention and are useful in a contemplated process. Inhibition data for those compounds in *in vitro* assays against MMP-1 and MMP-13 are provided
20 hereinafter.


Table 1




SUBSTITUTE SHEET (RULE 26)

Table 2


Example 9g




Example 13c




Example 10d




Example 11g




Example 14c




Example 12c



Example 15b



Example 16c



SUBSTITUTE SHEET (RULE 26)

Table 3













Table 4



SUBSTITUTE SHEET (RULE 26)


Table 5




SUBSTITUTE SHEET (RULE 26)

Table 6


Example 41




Example 42




Example 43




Example 44




Example 45



Example 46



Example 47



Example 48













Table 7









SUBSTITUTE SHEET (RULE 26)

Table 8



SUBSTITUTE SHEET (RULE 26)

Table 9



As utilized herein, the term "alkyl", alone or in combination, means a straight-chain or branched-chain alkyl radical containing 1 to about 12 carbon atoms, preferably 1 to about 10 carbon atoms, and more preferably 1 to about 6 carbon atoms. Examples of such radicals include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, iso-amyl, hexyl, octyl and the like.

The term "alkenyl", alone or in combination, means a straight-chain or branched-chain hydrocarbon radical having one or more double bonds and containing 2 to about 12 carbon atoms preferably 2 to about 10 carbon atoms, and more preferably, 2 to about 6 carbon atoms. Examples of suitable alkenyl radicals include ethenyl (vinyl), 2-propenyl, 3-propenyl, 1,4-pentadienyl, 1,4-butadienyl, 1-butenyl, 2-butenyl, 3-butenyl, decenyl and the like.

The term "alkynyl", alone or in combination, means a straight-chain hydrocarbon radical having one or more triple bonds and containing 2 to about 12 carbon atoms, preferably 2 to about 10 carbon atoms, and more preferably, 2 to about 6 carbon atoms. Examples of alkynyl radicals include ethynyl, 2-propynyl, 3-propynyl, decynyl, 1-butyne, 2-butyne, 3-butyne, and the like.

The term "carbonyl" or "oxo", alone or in combination, means a $-C(=O)-$ group wherein the remaining two bonds (valences) can be independently substituted. The term carbonyl is also intended to encompass a hydrated carbonyl group $-C(OH)_2-$.

The term "thiol" or "sulfhydryl", alone or in combination, means a $-SH$ group. The term "thio"

or "thia", alone or in combination, means a thiaether group; i.e., an ether group wherein the ether oxygen is replaced by a sulfur atom.

The term "amino", alone or in combination,
5 means an amine or $-NH_2$ group whereas the term mono-substituted amino, alone or in combination, means a substituted amine $-N(H)(\text{substituent})$ group wherein one hydrogen atom is replaced with a substituent, and disubstituted amine means a $-N(\text{substituent})_2$ wherein
10 two hydrogen atoms of the amino group are replaced with independently selected substituent groups.

Amines, amino groups and amides are compounds that can be designated as primary (I°), secondary (II°) or tertiary (III°) or unsubstituted,
15 mono-substituted or N,N-disubstituted depending on the degree of substitution of the amino nitrogen. Quaternary amine (ammonium) (IV°) means a nitrogen with four substituents $[-N^+(\text{substituent})_4]$ that is positively charged and accompanied by a counter ion,
20 whereas N-oxide means one substituent is oxygen and the group is represented as $[-N^+(\text{substituent})_3-O^-]$; i.e., the charges are internally compensated.

The term "cyano", alone or in combination, means a $-C$ -triple bond- N ($-C\equiv N$) group. The term
25 "azido", alone or in combination, means a $-N$ -triple bond- N ($-N\equiv N$) group. The term "hydroxyl", alone or in combination, means a $-OH$ group. The term "nitro", alone or in combination, means a $-NO_2$ group. The term "azo", alone or in combination, means a $-N=N-$
30 group wherein the bonds at the terminal positions can be independently substituted.

The term "hydrazino", alone or in combination, means a $-NH-NH-$ group wherein the

-52-

depicted remaining two bonds (valences) can be independently substituted. The hydrogen atoms of the hydrazino group can be replaced, independently, with substituents and the nitrogen atoms can form acid addition salts or be quaternized.

The term "sulfonyl", alone or in combination, means a $-SO_2-$ group wherein the depicted remaining two bonds (valences) can be independently substituted. The term "sulfoxido", alone or in combination, means a $-SO-$ group wherein the remaining two bonds (valences) can be independently substituted.

The term "sulfone", alone or in combination, means a $-SO_2-$ group wherein the depicted remaining two bonds (valences) can be independently substituted. The term "sulfenamide", alone or in combination, means a $-SON=$ group wherein the remaining three depicted bonds (valences) can be independently substituted. The term "sulfide", alone or in combination, means a $-S-$ group wherein the remaining two bonds (valences) can be independently substituted.

The term "alkoxy", alone or in combination, means an alkyl ether radical wherein the term alkyl is as defined above. Examples of suitable alkyl ether radicals include methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, iso-butoxy, sec-butoxy, tert-butoxy and the like.

The term "cycloalkyl", alone or in combination, means a cyclic alkyl radical that contains 3 to about 8 carbon atoms. The term "cycloalkylalkyl" means an alkyl radical as defined above that is substituted by a cycloalkyl radical containing 3 to about 8, preferably 3 to about 6,

SUBSTITUTE SHEET (RULE 26)

carbon atoms. Examples of such cycloalkyl radicals include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and the like.

A heterocyclic (heterocyclo) or heterocyclo
5 portion of a heterocyclocarbonyl, heterocyclooxy-
carbonyl, heterocycloalkoxycarbonyl, or
heterocycloalkyl group or the like is a saturated or
partially unsaturated monocyclic, bicyclic or
tricyclic heterocycle that contains one or more
10 hetero atoms selected from nitrogen, oxygen and
sulfur. Heterocyclo compounds include benzofused
heterocyclic compounds such as benzo-1,4-dioxane.
Such a moiety can be optionally substituted on one or
more ring carbon atoms by halogen, hydroxy,
15 hydroxycarbonyl, alkyl, alkoxy, oxo, and the like,
and/or on a secondary nitrogen atom (i.e., -NH-) of
the ring by alkyl, aralkoxycarbonyl, alkanoyl, aryl
or arylalkyl or on a tertiary nitrogen atom (i.e.,
=N-) by oxido and that is attached via a carbon atom.
20 The tertiary nitrogen atom with three substituents
can also attached to form a N-oxide [=N(O)-] group.

The term "aryl", alone or in combination,
means a 5- or 6-membered carbocyclic aromatic ring-
containing moiety or a fused ring system containing
25 two or three rings that have all carbon atoms in the
ring; i.e., a carbocyclic aryl radical. Exemplary
carbocyclic aryl radicals include phenyl, indenyl and
naphthyl radicals.

The term "heteroaryl", alone or in
30 combination means a 5- or 6-membered aromatic ring-
containing moiety or a fused ring system (radical)
containing two or three rings that have carbon atoms
and also one or more heteroatoms in the ring(s) such
as sulfur, oxygen and nitrogen. Examples of such

heterocyclic or heteroaryl groups are pyrrolidinyl, piperidyl, piperazinyl, morpholinyl, thiamorpholinyl, pyrrolyl, imidazolyl (e.g., imidazol-4-yl, 1-benzyloxycarbonylimidazol-4-yl, and the like),
5 pyrazolyl, pyridyl, pyrazinyl, pyrimidinyl, furyl, tetrahydrofuryl, thienyl, triazolyl, oxazolyl, oxadiazoyl, thiazolyl, thiadiazoyl, indolyl (e.g., 2-indolyl, and the like), quinolinyl, (e.g., 2-quinolinyl, 3-quinolinyl, 1-oxido-2-quinolinyl, and
10 the like), isoquinolinyl (e.g., 1-isoquinolinyl, 3-isoquinolinyl, and the like), tetrahydroquinolinyl (e.g., 1,2,3,4-tetrahydro-2-quinolyl, and the like), 1,2,3,4-tetrahydroisoquinolinyl (e.g., 1,2,3,4-tetrahydro-1-oxo-isoquinolinyl, and the like),
15 quinoxalinyl, β -carbolinyl, 2-benzofurancarbonyl, benzothiophenyl, 1-, 2-, 4- or 5-benzimidazolyl, and the like radicals.

When an aryl or heteroaryl radical is a substituting moiety (group, substituent, or radical),
20 it can itself substituted, the last-named substituent is independently selected from the group consisting of a cyano, perfluoroalkyl, trifluoromethoxy, trifluoromethylthio, haloalkyl, trifluoromethylalkyl, aralkoxycarbonyl,
25 aryloxycarbonyl, hydroxy, halo, alkyl, alkoxy, nitro, thiol, hydroxycarbonyl, aryloxy, arylthio, aralkyl, aryl, arylcarbonylamino, heteroaryloxy, heteroarylthio, heteroaralkyl, cycloalkyl, heterocyclooxy, heterocyclothio, heterocycloamino,
30 cycloalkyloxy, cycloalkylthio, heteroaralkoxy, heteroaralkylthio, aralkoxy, aralkylthio, aralkylamino, heterocyclo, heteroaryl, arylazo, hydroxycarbonylalkoxy, alkoxycarbonylalkoxy, alkanoyl, arylcarbonyl, aralkanoyl, alkanoyloxy,

aralkanoyloxy, hydroxyalkyl, hydroxyalkoxy,
alkylthio, alkoxyalkylthio, alkoxycarbonyl,
aryloxyalkoxyaryl, arylthioalkylthioaryl,
aryloxyalkylthioaryl, arylthioalkoxyaryl,
5 hydroxycarbonylalkoxy, hydroxycarbonylalkylthio,
alkoxycarbonylalkoxy, alkoxycarbonylalkylthio, amino,
wherein the amino nitrogen is (i) unsubstituted,
or (ii) substituted with one or two substituents
that are independently selected from the group
10 consisting of an alkyl, aryl, heteroaryl,
aralkyl, cycloalkyl, aralkoxycarbonyl,
alkoxycarbonyl, arylcarbonyl, aralkanoyl,
heteroarylcarbonyl, heteroaralkanoyl and an
alkanoyl group, or (iii) wherein the amino
15 nitrogen and two substituents attached thereto
form a 5- to 8-membered heterocyclo or
heteroaryl ring containing zero to two
additional heteroatoms that are nitrogen, oxygen
or sulfur and which ring itself is (a)
20 unsubstituted or (b) substituted with one or two
groups independently selected from the group
consisting of an aryl, alkyl, heteroaryl,
aralkyl, heteroaralkyl, hydroxy, alkoxy,
alkanoyl, cycloalkyl, heterocycloalkyl,
25 alkoxycarbonyl, hydroxyalkyl, trifluoromethyl,
benzofused heterocycloalkyl, hydroxyalkoxyalkyl,
aralkoxycarbonyl, hydroxycarbonyl,
aryloxycarbonyl, benzofused heterocycloalkoxy,
benzofused cycloalkylcarbonyl, heterocyclo-
30 alkylcarbonyl, and a cycloalkylcarbonyl group,
carbonylamino
wherein the carbonylamino nitrogen is (i)
unsubstituted, or (ii) is the reacted amine of
an amino acid, or (iii) substituted with one or

SUBSTITUTE SHEET (RULE 26)

-56-

two radicals selected from the group consisting of an alkyl, hydroxyalkyl, hydroxyheteroaralkyl, cycloalkyl, aralkyl, trifluoromethylalkyl, heterocycloalkyl, benzofused heterocycloalkyl, benzofused heterocycloalkyl, benzofused cycloalkyl, and an N,N-dialkylsubstituted alkylamino-alkyl group, or (iv) the carboxamido nitrogen and two substituents bonded thereto together form a 5- to 8-membered heterocyclo, heteroaryl or benzofused heterocycloalkyl ring that is itself unsubstituted or substituted with one or two radicals independently selected from the group consisting of an alkyl, alkoxycarbonyl, nitro, heterocycloalkyl, hydroxy, hydroxycarbonyl, aryl, aralkyl, heteroaralkyl and an amino group, wherein the amino nitrogen is (i) unsubstituted, or (ii) substituted with one or two substituents that are independently selected from the group consisting of alkyl, aryl, and heteroaryl, or (iii) wherein the amino nitrogen and two substituents attached thereto form a 5- to 8-membered heterocyclo or heteroaryl ring, and an aminoalkyl group wherein the aminoalkyl nitrogen is (i) unsubstituted, or (ii) substituted with one or two substituents independently selected from the group consisting of an alkyl, aryl, aralkyl, cycloalkyl, aralkoxycarbonyl, alkoxycarbonyl, and an alkanoyl group, or (iii) wherein the aminoalkyl nitrogen and two substituents attached thereto form a 5- to 8-membered heterocyclo or heteroaryl ring.

SUBSTITUTE SHEET (RULE 26)

The term "aralkyl", alone or in combination, means an alkyl radical as defined above in which one hydrogen atom is replaced by an aryl radical as defined above, such as benzyl, 2-phenylethyl and the like.

The term "aralkoxycarbonyl", alone or in combination, means a radical of the formula aralkyl-O-C(O)- in which the term "aralkyl" has the significance given above. An example of an aralkoxycarbonyl radical is benzyloxycarbonyl.

The term "aryloxy" means a radical of the formula aryl-O- in which the term aryl has the significance given above. The phenoxy radical is an exemplary aryloxy radical.

The terms "heteroaralkyl" and "heteroaryloxy" mean radicals structurally similar to aralkyl and aryloxy that are formed from heteroaryl radicals. Exemplary radicals include 4-picolinyl and 2-pyrimidinoxy, respectively.

The terms "alkanoyl" or "alkylcarbonyl", alone or in combination, means an acyl radical derived from an alkanecarboxylic acid, examples of which include formyl, acetyl, propionyl, butyryl, valeryl, 4-methylvaleryl, and the like.

The term "cycloalkylcarbonyl" means an acyl group derived from a monocyclic or bridged cycloalkanecarboxylic acid such as cyclopropanecarbonyl, cyclohexanecarbonyl, adamantanecarbonyl, and the like, or from a benzo-fused monocyclic cycloalkanecarboxylic acid that is optionally substituted by, for example, alkanoylamino, such as 1,2,3,4-tetrahydro-2-naphthoyl, 2-acetamido-1,2,3,4-tetrahydro-2-naphthoyl.

The terms "aralkanoyl" or "aralkylcarbonyl" mean an acyl radical derived from an aryl-substituted alkanecarboxylic acid such as phenylacetyl, 3-phenylpropionyl (hydrocinnamoyl), 4-phenylbutyryl, 5 (2-naphthyl)acetyl, 4-chlorohydrocinnamoyl, 4-aminohydrocinnamoyl, 4-methoxyhydrocinnamoyl and the like.

The terms "aroyl" or "arylcarbonyl" means an acyl radical derived from an aromatic carboxylic 10 acid. Examples of such radicals include aromatic carboxylic acids, an optionally substituted benzoic or naphthoic acid such as benzoyl, 4-chlorobenzoyl, 4-carboxybenzoyl, 4-(benzyloxycarbonyl)benzoyl, 1-naphthoyl, 2-naphthoyl, 6-carboxy-2 naphthoyl, 15 6-(benzyloxycarbonyl)-2-naphthoyl, 3-benzyloxy-2-naphthoyl, 3-hydroxy-2-naphthoyl, 3-(benzyloxyformamido)-2-naphthoyl, and the like.

The term "cycloalkylalkoxycarbonyl" means an acyl group of the formula cycloalkylalkyl-O-CO- 20 wherein cycloalkylalkyl has the significance given above. The term "aryloxyalkanoyl" means an acyl radical of the formula aryl-O-alkanoyl wherein aryl and alkanoyl have the significance given above. The term "heterocyclooxycarbonyl" means an acyl group 25 having the formula heterocyclo-O-CO- wherein heterocyclo is as defined above.

The term "heterocycloalkanoyl" is an acyl radical of the formula heterocyclo-substituted alkane carboxylic acid wherein heterocyclo has the 30 significance given above. The term "heterocycloalkoxycarbonyl" means an acyl radical of the formula heterocyclo-substituted alkane-O-CO- wherein heterocyclo has the significance given above. The term "heteroaryloxy carbonyl" means an acyl

radical represented by the formula heteroaryl-O-CO-
wherein heteroaryl has the significance given above.

The term "aminocarbonyl" (carboxamide)
alone or in combination, means an amino-substituted
5 carbonyl (carbamoyl) group derived from an amine
reacted with a carboxylic acid wherein the amino
(amido nitrogen) group is unsubstituted (-NH₂) or a
substituted primary or secondary amino group
containing one or two substituents selected from the
10 group consisting of hydrogen, alkyl, aryl, aralkyl,
cycloalkyl, cycloalkylalkyl radicals and the like, as
recited. A hydroxamate is a N-hydroxycarboxamide.

The term "aminoalkanoyl" means an acyl
group derived from an amino-substituted
15 alkanecarboxylic acid wherein the amino group can be
a primary or secondary amino group containing
substituents independently selected from hydrogen,
alkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl
radicals and the like.

20 The term "halogen" or "halo" means
fluoride, chloride, bromide or iodide. The term
"haloalkyl" means an alkyl radical having the
significance as defined above wherein one or more
hydrogens are replaced with a halogen. Examples of
25 such haloalkyl radicals include chloromethyl, 1-
bromoethyl, fluoromethyl, difluoromethyl,
trifluoromethyl, 1,1,1-trifluoroethyl and the like.

The term "perfluoroalkyl" means an alkyl
group wherein each hydrogen has been replaced by a
30 fluorine atom. Examples of such perfluoroalkyl
groups, in addition to trifluoromethyl above, are
perfluorobutyl, perfluoroisopropyl, perfluorododecyl
and perfluorodecyl.

The term "perfluoroalkoxy" alone or in combination, means a perfluoroalkyl ether radical wherein the term perfluoroalkyl is as defined above. Examples of such perfluoroalkoxy groups, in addition
5 to trifluoromethoxy ($\text{F}_3\text{C-O-}$), are perfluorobutoxy, perfluoroisopropoxy, perfluorododecyloxy and perfluorodecyloxy.

The term "perfluoroalkylthio" alone or in combination, means a perfluoroalkyl thioether radical
10 wherein the term perfluoroalkyl is as defined above. Examples of such perfluoroalkylthio groups, in addition to trifluoromethylthio ($\text{F}_3\text{C-S-}$), are perfluorobutylthio, perfluoroisopropylthio, perfluorododecylthio and perfluorodecylthio.

15 The term "aromatic ring" in combinations such as substituted-aromatic ring sulfone or substituted-aromatic ring sulfoxide means aryl or heteroaryl as defined before.


The term "pharmaceutically acceptable" is
20 used adjectivally herein to mean that the modified noun is appropriate for use in a pharmaceutical product. Pharmaceutically acceptable cations include metallic ions and organic ions. More preferred metallic ions include, but are not limited to
25 appropriate alkali metal (Group Ia) salts, alkaline earth metal (Group IIa) salts and other physiological acceptable metal ions. Exemplary ions include aluminum, calcium, lithium, magnesium, potassium, sodium and zinc in their usual valences. Preferred
30 organic ions include protonated tertiary amines and quaternary ammonium cations, including in part, trimethylamine, diethylamine, N,N'-dibenzylethylenediamine, chloroprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-

-61-

methylglucamine) and procaine. Exemplary
 pharmaceutically acceptable acids include without
 limitation hydrochloric acid, hydrobromic acid,
 phosphoric acid, sulfuric acid, methanesulfonic acid,
 5 acetic acid, formic acid, tartaric acid, maleic acid,
 malic acid, citric acid, isocitric acid, succinic
 acid, lactic acid, gluconic acid, glucuronic acid,
 pyruvic acid oxalacetic acid, fumaric acid, propionic
 acid, aspartic acid, glutamic acid, benzoic acid, and
 10 the like.

Treatment Process

A process for treating a host mammal having
 a condition associated with pathological matrix
 15 metalloprotease activity is also contemplated. That
 process comprises administering a compound
 corresponding in structure to Formula B that
 selectively inhibits one or more MMPs, while
 exhibiting less activity against at least MMP-1, in
 20 an MMP enzyme-inhibiting effective amount to a
 mammalian host having such a condition. A
 contemplated compound also does not substantially
 inhibit the production of TNF. The use of
 administration repeated a plurality of times is
 25 particularly contemplated.




B

wherein R¹, R², R⁶, R⁷, R¹³, and R¹⁴ are as
 defined before.

SUBSTITUTE SHEET (RULE 26)

-62-

A more preferred group of contemplated compounds for use in such a process corresponds in structure to Formula B1, below,



B1

5

wherein R¹ and R¹⁴ are as defined before,
and is a substituent selected from the group
consisting of an alkylcarbonyl, alkoxycarbonyl,
10 arylcarbonyl, alkylsulfonyl, arylsulfonyl,
aminoalkylcarbonyl, aminoaralkylcarbonyl,
alkoxycarbonylaminoalkylcarbonyl,
alkoxycarbonylaminoaralkylcarbonyl,
cycloalkoxycarbonylaminoalkylcarbonyl,
15 cycloalkoxycarbonylaminoaralkylcarbonyl,
aryloxycarbonylaminoalkylcarbonyl, aryloxycarbonyl-
aminoaralkylcarbonyl, aralkyloxycarbonylamino-
alkylcarbonyl, aralkyloxycarbonylaminoalkylcarbonyl,
heteroaryloxycarbonylaminoalkylcarbonyl and a
20 heteroaryloxycarbonylaminoaralkylcarbonyl group.

A contemplated compound is used for
treating a host mammal such as a mouse, rat, rabbit,
dog, horse, primate such as a monkey, chimpanzee or
human that has a condition associated with
25 pathological matrix metalloprotease activity.

Also contemplated is the similar use of a
contemplated compound in the treatment of a disease
state that can be affected by the activity of
metalloproteases such as TNF- α convertase. Exemplary

SUBSTITUTE SHEET (RULE 26)

of such disease states are the acute phase responses of shock and sepsis, coagulation responses, hemorrhage and cardiovascular effects, fever and inflammation, anorexia and cachexia.

5 In treating a disease condition associated with pathological matrix metalloproteinase activity, a contemplated MMP inhibitor compound can be used, where appropriate, in the form of an amine salt derived from an inorganic or organic acid. Exemplary
10 acid salts include but are not limited to the following: acetate, adipate, alginate, citrate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, camphorate, camphorsulfonate, digluconate, cyclopentanepropionate, dodecylsulfate,
15 ethanesulfonate, glucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, fumarate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxy-ethanesulfonate, lactate, maleate, methanesulfonate, nicotinate, 2-naphthalenesulfonate, oxalate,
20 palmoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, tosylate, mesylate and undecanoate.

Also, a basic nitrogen-containing group can be quaternized with such agents as lower alkyl (C_1 -
25 C_6) halides, such as methyl, ethyl, propyl, and butyl chloride, bromides, and iodides; dialkyl sulfates like dimethyl, diethyl, dibutyl, and diamyl sulfates, long chain (C_8 - C_{20}) halides such as decyl, lauryl, myristyl and dodecyl chlorides, bromides and iodides,
30 aralkyl halides like benzyl and phenethyl bromides, and others to provide enhanced water-solubility. Water or oil-soluble or dispersible products are thereby obtained as desired. The salts are formed by combining the basic compounds with the desired acid.

SUBSTITUTE SHEET (RULE 26)

Other compounds useful in this invention that are acids can also form salts. Examples include salts with alkali metals or alkaline earth metals, such as sodium, potassium, calcium or magnesium or
5 with organic bases or basic quaternary ammonium salts.

In some cases, the salts can also be used as an aid in the isolation, purification or resolution of the compounds of this invention.

10 Total daily dose administered to a host mammal in single or divided doses of an MMP enzyme-inhibiting effective amount can be in amounts, for example, of about 0.001 to about 100 mg/kg body weight daily, preferably about 0.001 to about 30
15 mg/kg body weight daily and more usually about 0.01 to about 10 mg. Dosage unit compositions can contain such amounts or submultiples thereof to make up the daily dose. A suitable dose can be administered, in multiple sub-doses per day. Multiple doses per day
20 can also increase the total daily dose, should such dosing be desired by the person prescribing the drug.

The dosage regimen for treating a disease condition with a compound and/or composition of this invention is selected in accordance with a variety of
25 factors, including the type, age, weight, sex, diet and medical condition of the patient, the severity of the disease, the route of administration, pharmacological considerations such as the activity, efficacy, pharmacokinetic and toxicology profiles of
30 the particular compound employed, whether a drug delivery system is utilized and whether the compound is administered as part of a drug combination. Thus, the dosage regimen actually employed can vary widely

-65-

and therefore can deviate from the preferred dosage regimen set forth above.

A compound useful in the present invention can be formulated as a pharmaceutical composition.

- 5 Such a composition can then be administered orally, parenterally, by inhalation spray, rectally, or topically in dosage unit formulations containing conventional nontoxic pharmaceutically acceptable carriers, adjuvants, and vehicles as desired.
- 10 Topical administration can also involve the use of transdermal administration such as transdermal patches or iontophoresis devices. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal
- 15 injection, or infusion techniques. Formulation of drugs is discussed in, for example, Hoover, John E., Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, Pennsylvania; 1975 and Liberman, H.A. and Lachman, L., Eds., Pharmaceutical Dosage Forms, Marcel Decker, New York, N.Y., 1980.
- 20

- Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions can be formulated according to the known art using suitable dispersing or wetting agents and suspending
- 25 agents. The sterile injectable preparation can also be a sterile injectable solution or suspension in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that can be
- 30 employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono-

SUBSTITUTE SHEET (RULE 26)

-66-

or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables. Dimethyl acetamide, surfactants including ionic and non-ionic detergents, polyethylene glycols can be used. Mixtures of solvents and wetting agents such as those discussed above are also useful.

Suppositories for rectal administration of the drug can be prepared by mixing the drug with a suitable nonirritating excipient such as cocoa butter, synthetic mono- di- or triglycerides, fatty acids and polyethylene glycols that are sold at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum and release the drug.

Solid dosage forms for oral administration can include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the compounds of this invention are ordinarily combined with one or more adjuvants appropriate to the indicated route of administration. If administered per os, the compounds can be admixed with lactose, sucrose, starch powder, cellulose esters of alcanoic acids, cellulose alkyl esters, talc, stearic acid, magnesium stearate, magnesium oxide, sodium and calcium salts of phosphoric and sulfuric acids, gelatin, acacia gum, sodium alginate, polyvinylpyrrolidone, and/or polyvinyl alcohol, and then tableted or encapsulated for convenient administration. Such capsules or tablets can contain a controlled-release formulation as can be provided in a dispersion of active compound in hydroxypropylmethyl cellulose. In the case of capsules, tablets, and pills, the dosage forms can also comprise buffering agents such as sodium

SUBSTITUTE SHEET (RULE 26)

citrate, magnesium or calcium carbonate or bicarbonate. Tablets and pills can additionally be prepared with enteric coatings.

For therapeutic purposes, formulations for
5 parenteral administration can be in the form of aqueous or non-aqueous isotonic sterile injection solutions or suspensions. These solutions and suspensions can be prepared from sterile powders or granules having one or more of the carriers or
10 diluents mentioned for use in the formulations for oral administration. The compounds can be dissolved in water, polyethylene glycol, propylene glycol, ethanol, corn oil, cottonseed oil, peanut oil, sesame oil, benzyl alcohol, sodium chloride, and/or various
15 buffers. Other adjuvants and modes of administration are well and widely known in the pharmaceutical art.

Liquid dosage forms for oral administration can include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs
20 containing inert diluents commonly used in the art, such as water. Such compositions can also comprise adjuvants, such as wetting agents, emulsifying and suspending agents, and sweetening, flavoring, and perfuming agents.

25 The amount of active ingredient that can be combined with the carrier materials to produce a single dosage form varies depending upon the mammalian host treated and the particular mode of administration.

30

Compound Preparation

A contemplated compound can be prepared by several routes of synthesis as can be seen from the specific Examples that follow. In addition, Schemes

1A and 1B, below, illustrate exemplary synthetic procedures and the use of preferred intermediates in the preparation of contemplated compounds.


Thus, as is seen in Scheme 1A, methyl N-
5 acetylacrylate, compound 1, is reacted in a solvent such as methanol with p-fluorothiophenol in the presence of triethylamine at zero degrees C via a Michael addition (step 1) and the resulting fluorophenylthioether is oxidized with Oxone® to form
10 the corresponding sulfone, compound 2, in step 2. Hydrolysis of the methyl ester in glacial acetic acid containing hydrochloric acid at zero degrees C provides the corresponding acid, compound 3. Reaction of compound 3 with thiophenol in DMF at
15 about 10° C in the presence of potassium carbonate provides the thiophenylether, compound 4. The N-acetyl group is cleaved by reaction of compound 4 in glacial acetic acid and hydrochloric acid at 11° C to provide the amine hydrochloride, compound 5.

20 Reaction of compound 5 in water with an N-protected amino acid such as CBZ-glycine (CBZ-Gly) in the presence of a water-soluble carbodiimide [1-ethyl-3-[3-(dimethylamino)-propyl]carbodiimide hydrochloride; EDC], 1-hydroxybenzotriazole (HOBT)
25 and N-methylmorpholine (NMM) provides the corresponding N-aryloxycarbonylaminoalkylcarbonyl derivative, compound 6, that is shown in Scheme 1B. Reaction of compound 6 with O-tetrahydro-2H-pyran-2-yl-hydroxylamine in THF in the presence of HOBT and
30 EDC provides the THP-protected hydroxamic acid, Compound 7. Reaction of compound 7 with an acid such as toluenesulfonic acid or hydrochloric acid cleaves the THP protecting group to provide the hydroxamic acid, compound 8.


SUBSTITUTE SHEET (RULE 26)

- 69 -

Scheme 1A



Scheme 1B



SUBSTITUTE SHEET (RULE 26)

The above syntheses, as with all of the reactions discussed herein, can be carried out under a dry inert atmosphere such as nitrogen, helium or argon if desired. Selected reactions known to those skilled in the art can be carried out under a dry atmosphere such as dry air, whereas other synthetic steps, for example, aqueous acid or base ester or amide hydrolyses, can be carried out under laboratory air.

10 In general, the choices of starting material and reaction conditions can vary as is well known to those skilled in the art. Usually, no single set of conditions is limiting because variations can be applied as required and selected by one skilled in the art. Conditions can also be selected as desired to suit a specific purpose such as small scale preparations or large scale preparations. In either case, the use of less safe or less environmentally sound materials or reagents is usually to be minimized. Examples of such less desirable materials are diazomethane, diethyl ether, heavy metal salts, dimethyl sulfide, chloroform, benzene and the like.

25 Various reactions illustrated in the above Scheme can be base-mediated by the use of catalytic amounts of some bases or carried out with an equivalent or more of a base by the addition of an additional reagent or the thiol reagent can be a preformed thiol salt such as the sodium salt of a thiophenol. Bases that can be used include, for example, metal hydroxides such as sodium, potassium, lithium or magnesium hydroxide, oxides such as those of sodium, potassium, lithium, calcium or magnesium, metal carbonates such as those of sodium, potassium,

lithium, calcium or magnesium, metal bicarbonates such as sodium bicarbonate or potassium bicarbonate, primary (I°), secondary (II°) or tertiary (III°) organic amines such as alkyl amines, arylalkyl
5 amines, alkylarylalkyl amines, heterocyclic amines or heteroaryl amines, ammonium hydroxides or quaternary ammonium hydroxides.

As non-limiting examples, such amines can include triethyl amine, trimethylamine, diisopropyl-
10 amine, methyldiisopropylamine, diazabicyclononane, tribenzylamine, dimethylbenzylamine, morpholine, N-methylmorpholine, N,N'-dimethylpiperazine, N-ethylpiperidine, 1,1,5,5-tetramethylpiperidine, dimethylaminopyridine, pyridine, quinoline,
15 tetramethylethylenediamine and the like. Non-limiting examples of ammonium hydroxides, usually made from amines and water, can include ammonium hydroxide, triethylammonium hydroxide, trimethyl-
ammonium hydroxide, methyldiisopropylammonium
20 hydroxide, tribenzylammonium hydroxide, dimethylbenzyl-ammonium hydroxide, morpholinium hydroxide, N-methylmorpholinium hydroxide, N,N'-dimethylpiperazinium hydroxide, N-ethylpiperidinium hydroxide, and the like. As non-limiting examples,
25 quaternary ammonium hydroxides can include tetraethyl-ammonium hydroxide, tetramethylammonium hydroxide, dimethyldiisopropylammonium hydroxide, benzylmethyldiisopropylammonium hydroxide, methyldiazabicyclononylammonium hydroxide,
30 methyltribenzylammonium hydroxide, N,N'-dimethylmorpholiniumhydroxide, N,N,N', N',-tetramethylpiperaziniumhydroxide, and N-ethyl-N'-hexylpiperidinium hydroxide and the like.

Metal hydrides, amides or alcoholates such as calcium hydride, sodium hydride, potassium hydride, lithium hydride, sodium methoxide, potassium tert-butoxide, calcium ethoxide, magnesium ethoxide, sodium amide, potassium diisopropyl amide and the like may also be suitable reagents. Organometallic deprotonating agents such as alkyl or aryl lithium reagents such as methyl, phenyl or butyl lithium, Grignard reagents such as methylmagnesium bromide or methymagnesium chloride, organocadium reagents such as dimethylcadium and the like can also serve as bases for causing salt formation or catalyzing the reaction. Quaternary ammonium hydroxides or mixed salts are also useful for aiding phase transfer couplings or serving as phase transfer reagents.

The reaction media can comprise a single solvent, mixed solvents of the same or different classes or serve as a reagent in a single or mixed solvent system. The solvents can be protic, non-protic or dipolar aprotic. Non-limiting examples of protic solvents include water, methanol (MeOH), denatured or pure 95% or absolute ethanol, isopropanol and the like. Typical non-protic solvents include acetone, tetrahydrofuran (THF), dioxane, diethylether, tert-butylmethyl ether (TBME), aromatics such as xylene, toluene, or benzene, ethyl acetate, methyl acetate, butyl acetate, trichloroethane, methylene chloride, ethylenedichloride, hexane, heptane, isooctane, cyclohexane and the like. Dipolar aprotic solvents include compounds such as dimethylformamide (DMF), dimethylacetamide (DMAc), acetonitrile, nitromethane, tetramethylurea, N-methylpyrrolidone and the like.

Non-limiting examples of reagents that can be used as solvents or as part of a mixed solvent system include organic or inorganic mono- or multi-protic acids or bases such as hydrochloric acid, phosphoric acid, sulfuric acid, acetic acid, formic acid, citric acid, succinic acid, triethylamine, morpholine, N-methylmorpholine, piperidine, pyrazine, piperazine, pyridine, potassium hydroxide, sodium hydroxide, alcohols or amines for making esters or amides or thiols for making the products of this invention and the like.

Room temperature or less or moderate warming (-10°C to 60°C) are the preferred temperatures of the reaction. If desired, the reaction temperature can be about -76°C to the reflux point of the reaction solvent or solvents.

An intermediate thioether such as the 4-fluorophenylthio ether can be oxidized to the sulfone in one step using two equivalents to oxidizing agent. Reagents for this process can, in a non-limiting example, include peroxymonosulfate (OXONE[®]), hydrogen peroxide, meta-chloroperbenzoic acid, perbenzoic acid, peracetic acid, perlactic acid, tert-butyl peroxide, tert-butyl hydroperoxide, tert-butyl hypochlorite, sodium hypochlorite, hypochlorous acid, sodium meta-periodate, periodic acid and the like. Protic, non-protic, dipolar aprotic solvents, either pure or mixed, can be chosen, for example, methanol/water.

The oxidation can be carried out at temperature of about -78° to about 50° degrees centigrade, and normally is in a range of about -10°C to about 40°C . Preparation of a desired sulfone can be carried out in a two-step process using about one

equivalent of oxidizing agent to first form the sulfoxide at about 0°C. A second oxidation then produces the sulfone.

The solvents listed above can be used with these selective oxidations with, for example, methanol or methanol/water being preferred along with a temperature of from about -10°C to 30°C. It can be desirable in the case of more active oxidizing agents, but not required, that the reactions be carried out under an inert gas atmosphere with or without degassed solvents.

A hydroxamate can be prepared from the corresponding ester by reaction of the ester with one or more equivalents of hydroxylamine hydrochloride at room temperature or above in a solvent or solvents such as those listed above. This exchange process can be further catalyzed by the addition of additional acid.

Alternatively, a base such as a salt of an alcohol used as a solvent, for example, sodium methoxide in methanol, can be used to form hydroxylamine in situ, which can exchange with an ester or amide. The exchange can be carried out with a protected hydroxylamine such as O-tetrahydro-2H-pyran-2-yl-hydroxylamine (tetrahydropyranyl-hydroxyamine; THPONH₂), benzylhydroxylamine (BnONH₂), and the like in which case compounds in which the ester is a tetrahydropyranyl (THP) or benzyl (Bn) ester.

Removal of the protecting groups when desired, for example, following further transformations in another part of the molecule or following storage, is accomplished by standard methods well known in the art such as acid hydrolysis

SUBSTITUTE SHEET (RULE 26)

of the THP group with an acid such as toluenesulfonic
or hydrochloric acids or reductive removal of the
benzyl group with hydrogen and a metal catalyst such
as palladium, platinum, palladium on carbon or
5 nickel.

Oxidizable functional groups are readily
recognized by those skilled in the art and
alternative synthesis can be used such as the
protection/deprotection sequence.

10 Acids can be converted into activated
carbonyl compounds using reagents well known in the
art including the peptide and protein synthesis and
amino acid coupling or conjugation art. Examples of
such reagents are thionyl chloride, oxalyl chloride,
15 phosphorus oxychloride, 1-hydroxybenzotriazole
(HOBT), isobutylchloroformate and the like. These
valuable activated carbonyl intermediates can then be
transformed into hydroxamic acids or hydroxamic acid
derivatives such as H, benzyl or THP. Preparation of
20 or interconversion between the hydroxylamine or
hydroxylamine derivative compounds or acids or amides
or esters can be carried out by one skilled in the
art using the methods discussed above or by other
techniques.

25 The amine function in the intermediate
compounds use a protecting group to facilitate the
transformations. Decisions involving the selection
of protecting groups and their use can be made by a
person skilled in the art. Especially useful are the
30 techniques and reagents used in protein, peptide and
amino acid coupling and transformation chemistry.
The use of the tert-butoxycarbonyl (BOC),
benzyloxycarbonyl (Z or CBZ) and N,N-dibenzyl groups

as well as their synthesis and removal are examples of such protection schemes.

Coupling of the amino acids, amino acid esters, amino acid hydroxamates or hydroxamate
5 derivatives and amino acid amides of the precursor (intermediate) compounds with, for example, other amino acids, amines, alcohols, amides or acids is also carried out by methods well known in the art such as, for example, active ester or mixed anhydride
10 couplings with preferred bases if required being moderate tertiary amines such as N-methylmorpholine. Removal of a preexisting group that can also serve as a protecting group or blocking group such as the acetyl group and the like is also accomplished using
15 standard hydrolysis conditions such as base hydrolysis or exchange or acid exchange or hydrolysis.

In the case of compounds with an amine group, it is sometimes desirable to use acidic
20 conditions with a reagent such as hydrogen peroxide and/or in combination with an acidic reagent such as periodic acid, peracetic acid and the like. It should also be noted by one skilled in the art that hydrolysis or exchange of the acetyl group may or may
25 not effect hydrolysis or exchange of an ester, amide or hydroxamate function.

A selectively removable protecting group, P (R^{22}), on the hydroxamate hydroxy group can also be utilized. Such protecting groups can include acyl
30 groups, carbamoyl groups, ethers, alkoxyalkyl ethers, cycloalkyloxy ethers, arylalkyl groups, trisubstituted silyl groups and the like. Examples of such protecting groups include acetyl, THP, benzyl, Z, tert-butyldimethylsilyl (TBDMS) groups.

- 78 -

The preparation of such protected hydroxamates as well as the removal of the protecting groups is well known in the art and its practitioners.

Compounds of this invention can be prepared
5 by alkylation of a carbanion (nucleophile) generated from a protected carboxylic acid using processes known in the art. Protecting groups for the carboxyl function include, for example, esters such as tert-butyl esters. Bases for forming the anion are can be
10 organometallic reagents such as tert-butyl lithium, metal amides such as lithium diisopropyl amide (LDA) or alkoxides such as potassium tert-butoxide. Other candidate bases are discussed above.

Following or during formation of the anion,
15 the alkylating agent (electrophile) is added which undergoes a nucleophilic substitution reaction. Electrophilic substrates for displacement can include, for example, dihalo alkanes such as 1,2-dihaloalkanes or mono-halo-mono sulfated alkanes or
20 bissulfonate alkane esters. 1,2-di-Bromoethanes, 1-chloro-2, bromoethanes, 1-chloro-2-tosylethanes and 1,2-di-toluenesulfonylethanes are examples of such bis-electrophiles. 1-Bromo-2-chloro-ethane is a preferred electrophile.

25 Activated ester groups are well known in the art and can include, for example, di-esters such as malonates, ester-ketones such as acetoacetic esters or ester-aldehydes that are subject to carbonyl addition reactions. Alkylation with one
30 equivalent of alkylating agent followed by derivatization of the new omega carbonyl group with, for example, an organometallic reagent or reduction to form an alcohol which can then be derivatized to form a carbon halogen bonds or an activated ester

SUBSTITUTE SHEET (RULE 26)

such as a sulfate ester. These omega-substituted compounds can serve as substrates for the thioate displacement and oxidation reactions discussed above to form the carboxylic acid compounds or
5 intermediates of this invention.

Omega-haloalcohols can be useful starting materials for the preparation of compounds of this invention using alternative synthetic sequences from those discussed above. They can serve as substrates
10 for R^1 thiolate displacement (SN_2) to provide 4-sulfides (thio ethers) that can then be oxidized to the desired sulfones. The $HS-R^1$ compounds can be prepared as discussed below and oxidized as discussed above. Preparation of the R^1 group can be via an
15 intermediate such as a fluorothiophenol followed by displacement of the fluoride with a second nucleophile to produce compounds or intermediates of this invention. Fluorothiophenol and phenol and 2,3-dimethyl phenol are examples of preferred thiols and
20 phenols, respectively. The sulfone alcohols can be oxidized to the corresponding carboxylic acids as well as to the corresponding aldehydes.

The preparation of ethers can be carried out by forming a salt of the alcohol and treating
25 this nucleophile with an electrophile such as a halide or an activated ester such as a sulfate ester. The salt is formed by treating the alcohol with a base such as is discussed above. Examples of such bases are lithium alkyls, metal hydrides or the metal
30 salts of an amine such as LDA.

Halides can be chlorides, bromides or iodides and sulfates can be, for example, benzene sulfonates, tosylates, mesylates or triflates. An example of a preferred electrophile is 2-

SUBSTITUTE SHEET (RULE 26)

chloromethylpyridine and a preferred base is sodium
hydride. Alternatively, the alcohol can be converted
into a leaving group (electrophilic reagent) and then
treated with a nucleophile. Examples of such leaving
5 groups include sulfate esters such tosylates,
mesylates and triflates whose preparation is
discussed above. The triflate is a preferred leaving
group.

Displacement of these groups with
10 nucleophiles is well known in the art and discussed
and/or illustrated above. The nucleophiles can be
hydroxide to allow inversion of stereochemistry,
alkoxides to form ethers, amines or ammonia to form
substituted amines or an azide anion to form an
15 azide. A preferred nucleophile the is tetra-(n-
butyl)ammonium azide. The azido compound, for
example, can be reduced to form the amino acid.
Reductions are discussed above and are well known in
the art. A preferred method is hydrogenation with
20 palladium on carbon catalyst.

The amines, including the amino acids, of
this invention can be acylated or alkylated by
methods well known in the art. The amides formed can
be considered as protected amines or as end products
25 of this invention. Acylation to form such
derivatives as tert-butoxycarbonyl and carbobenzyloxy
carbonyl group is discussed above. Other acyl (Ac)
groups can be, for example, acetyl, haloacetyl,
aroyl, substituted aroyl, heteroaroyl, substituted
30 heteroaroyl or other groups as required. The amines
can be acylated using anhydrides, mixed anhydrides,
acid chlorides or activated esters. Usually such
acylations are carried out in presence of a base such
as the bases discussed above and well known in the

art. Examples are N-methyl-morpholine, triethylamine and the like.

The carboxyl compounds useful herein having amide substituents can be treated, converted or
5 interconverted as shown and/or discussed above to form the products of this invention. In addition, the haloacetyl compounds such as the preferred 2-chloroacetamide derivative can be treated with an amine as a nucleophile to yield an amino acid.
10 Again, these reactions are well known in the art. A preferred amine is morpholine.

Alkylation of the amino acid at the carbon alpha to the carbonyl group to form a useful compound can be carried out by first forming an anion using a
15 base. Exemplary bases are discussed elsewhere. The preferred bases are strong bases that are either hindered and/or non-nucleophilic such as lithium amides, metal hydrides or lithium alkyls. A preferred base is lithium diisopropylamide (LDA) in a
20 dipolar aprotic solvent or THF.

Following or during formation of the anion, an alkylating agent (an electrophile) is added which undergoes a nucleophilic substitution reaction. Non-limiting examples of such alkylating agents are 1,2-
25 dihaloalkanes or haloalkanes also substituted by an activated ester group. Activated ester groups are well known in the art and can include, for example, an ester of a 2-halo-alcohol such as a bromo-, iodo- or chloro-ethane para-toluene sulfonate, triflate or
30 mesylate. A preferred alkylating agents is 1-bromo-2-chloroethane.

The nitrogen substituent on the amino acid portion of the compounds of this invention can be varied. In addition, this can be accomplished at

different stages in the synthetic sequence based on the needs and objectives of the skilled person preparing the compounds of this invention.

The N-side chain variations can include

5 replacing the hydrogen (hydrido) substituent with a alkyl, arylalkyl, alkene or alkyne. This can be accomplished by methods well known in the art such as alkylation of the amine with an electrophile such as halo- or sulfate ester (activated ester) derivative

10 of the desired side chain. This can be done in the presence of a base such as those discussed above and in a pure or mixed solvent as discussed above. A preferred base is potassium carbonate and a preferred solvent is DMF.

15 The alkenes and alkynes can be reduced, if desired, by, for example, hydrogenation with a metal catalyst and hydrogen, to an alkyl or arylalkyl compound of this invention and the alkyne or arylalkyne can be reduced to a alkene of alkane with

20 under catalytic hydrogenation conditions as discussed above or with an deactivated metal catalyst. Catalysts can include, for example, Pd, Pd on Carbon, Pt, PtO₂ and the like. Less robust catalysts include such thing as Pd on BaCO₃ or Pd with quinoline or/and

25 sulfur.

An alternative method for alkylation of the amine nitrogen is reductive alkylation. This process, well known in the art, allows treatment of the secondary amine with an aldehyde or ketone in the

30 presence of a reducing agent such as borane, borane:THF, borane:pyridine, lithium aluminum hydride. Alternatively, reductive alkylation can be carried out hydrogenation conditions in the presence of a metal catalyst. Catalysts, hydrogen pressures

and temperatures are discussed above and are well known in the art. A preferred reductive alkylation catalyst is borane:pyridine complex.

The compounds of this invention include
5 compounds wherein the substituent on nitrogen of the amino acids as listed above provide amino acid carbamates. Non-limiting examples of these carbamates are the carbobenzoxy carbonyl (Z, CBZ, benzyloxycarbonyl), isobutoxycarbonyl and tert-
10 butoxycarbonyl (BOC, t-BOC) compounds. These materials can be made, as discussed above, at various stages in the synthesis based on the needs and decisions made by a person skilled in the art using methods well known in the art.

15 Useful synthetic techniques and reagents include those used in protein, peptide and amino acid synthesis, coupling and transformation chemistry. The use of the tert-butoxycarbonyl (BOC) and benzyloxycarbonyl (Z) as well as their synthesis and
20 removal are examples of such protection or synthesis schemes discussed above. Transformations of amino acids, amino esters, amino acid hydroxamates, amino acid hydroxamate derivatives and amino acid amides of this invention or compounds used in this invention
25 can be carried out as discussed and/or illustrated above. This includes, for example, active ester or mixed anhydride couplings wherein preferred bases, if required, are tertiary amines such as N-methylmorpholine.

30 Reagents for protection of the amine group of the protected amino acids include carbobenzoxy chloride, iso-butylchloroformate, tert-butoxycarbonyl chloride, di-tert-butyl dicarbonate and the like which are reacted with the amine in non-protic or

dipolar aprotic solvents such as DMF or THF or mixtures of solvents. A preferred reagent is di-tert-butyl dicarbonate and a preferred solvent is THF. Further conversion of the cyclic amino acids of this invention including alkylation, displacement with a thiol or thiolate, oxidation to a sulfone, and conversion into a hydroxamic acid or hydroxamate derivative can be carried out discussed herein.

Sulfone compounds such as those where R^1 is nitroaryl can be prepared as compounds of this invention by synthesis of a thiol or thiolate nucleophile, displacement of an electrophile (X) by the nucleophilic thiol or thiolate and oxidation of the product thio ether (sulfide) to the sulfone. For example, displacement of the electrophilic group X with a nitro-benzenethiol can yield a compound where R^1 is nitrobenzene that can be reduced to provide a useful amino compound wherein R^1 is an aniline that can be acylated to form a benzamido compound such as compounds 61-64 of Table 8, for example. It should be noted that nitrobenzenethiol is an example and not to be considered as limiting or required. Oxidation of the thioether product can be carried out as discussed below when desired.

The reduction of nitro groups to amines is well known in the art with a preferred method being hydrogenation. There is usually a metal catalyst such as Rh, Pd, Pt, Ni or the like with or without an additional support such as carbon, barium carbonate and the like. Solvents can be protic or non-protic pure solvents or mixed solvents as required. The reductions can be carried out at atmospheric pressure to a pressure of multiple atmospheres with atmospheric pressure to about 40 pounds per square

inch (psi) preferred. The amino group can be alkylated if desired, or acylated with, for example, an aroyl chloride, heteroaryl chloride or other amine carbonyl forming agent to form an R¹ amide.

5 The amino sulfone or thioether can also be reacted with a carbonic acid ester chloride, a sulfonyl chloride, a carbamoyl chloride or an isocyanate to produce the corresponding carbamate, sulfonamides, or urea. Acylation of amines of this
10 type are well known in the art and the reagents are also well known.

 Usually, these reactions are carried out in aprotic solvents under an inert or/and dry atmosphere at about 45°C to about -10°C. An equivalent of a
15 non-competitive base is usually used with sulfonyl chloride, acid chloride or carbonyl chloride reagents. Following or before this acylation step, synthesis of the hydroxamic acid products of this invention can proceed as discussed.

20 Other thiol reagents can also be used in the preparation of compounds of this invention. Examples are fluoroaryl, fluoroheteroaryl, azidoaryl or azidoheteroaryl or heteroaryl thiol reagents. These thiols can be used as nucleophiles to as
25 discussed above. Oxidation to the corresponding sulfone can then be carried out. The fluoro-substituted sulfone can be treated with a nucleophile such as ammonia, a primary amine, a quaternary ammonium or metal azide salt, under pressure if
30 desired, to provide an azido, amino or substituted amino group that can then be reacted with an activated benzoic or substituted benzoic acid derivative to form a benzamide. Azides can be reduced to an amino group using, for example, hydrogen with a metal

SUBSTITUTE SHEET (RULE 26)

catalyst or metal chelate catalyst or by an activated hydride transfer reagent. Hydrazo compounds can be oxidized to azo compounds and axo compounds can be reduced to hydrazo compounds. The amines can be
5 acylated as discussed above.

Preferred methods of preparing aminethiol intermediates of this invention include protection of an aromatic or heteroaromatic thiol with trityl chloride to form the trityl thiol derivative,
10 treatment of the amine with as reagent such as an aromatic or heteraromatic acid chloride to form the amide, removal of the trityl group, with acid to form the thiol. Preferred acylating agents include benzoyl chloride and preferred trityl removing
15 reagents include trifluoroacetic acid and trisopropylsilane.

The fluorine on fluorosulfone intermediates can also be displaced with other aryl or heteroaryl nucleophiles for form compounds of this invention.
20 Examples of such nucleophiles include salts of phenols, thiophenols, -OH group containing aromatic heterocyclic compounds or -SH containing heteroaryl compounds.

Tautomers of such groups azo, hydrazo, -OH
25 or -SH are specifically included as useful isomers. A preferred method of preparing intermediates in the synthesis of the substituted sulfones is by oxidation of an appropriate acetophenone, prepared from a fluoroacetophenone, with for example,
30 peroxymonosulfate, to form the corresponding phenol-ether. That phenol-ether is converted into its dimethylthiocarbamoyl derivative using dimethylthiocarbamoyl chloride, followed by rearranging the dimethylthiocarbamoyl derivative with

-87-

heat to provide the thiol required for preparation of the thioether intermediate.

Salts of the compounds or intermediates of this invention are prepared in the normal fashion
5 wherein acidic compounds are reacted with bases such as those discussed above to produce metal or nitrogen containing cation salts. Basic compounds such as amines can be treated with an acid to form the amine salt. A preferred amine salt is the
10 hydrochloride salt formed by reaction of the free base with HCl or hydrochloric acid.

Compounds of the present invention can possess one or more asymmetric carbon atoms and are thus capable of existing in the form of optical
15 isomers as well as in the form of racemic or non-racemic mixtures thereof. The optical isomers can be obtained by resolution of the racemic mixtures according to conventional processes well known in the art, for example by formation of diastereoisomeric
20 salts by treatment with an optically active acid or base. Examples of appropriate acids are tartaric, diacetyltartaric, dibenzoyltartaric, ditoluoyltartaric and camphorsulfonic acid and then separation of the mixture of diastereoisomers by
25 crystallization followed by liberation of the optically active bases from these salts. A different process for separation of optical isomers involves the use of a chiral chromatography column optimally chosen to maximize the separation of the enantiomers.

30 Still another available method involves synthesis of covalent diastereoisomeric molecules, e.g., esters, amides, acetals, ketals, and the like, by reacting compounds of Formula I with an optically active acid in an activated form, a optically active

SUBSTITUTE SHEET (RULE 26)

diol or an optically active isocyanate. The synthesized diastereoisomers can be separated by conventional means such as chromatography, distillation, crystallization or sublimation, and then hydrolyzed to deliver the enantiomerically pure compound. In some cases hydrolysis to the parent optically active drug is not necessary prior to dosing the patient because the compound can behave as a prodrug. The optically active compounds of Formula I can likewise be obtained by utilizing optically active starting materials.

In addition to the optical isomers or potentially optical isomers discussed above, other types of isomers are specifically intended to be included in this discussion and in this invention. Examples include cis isomers, trans isomers, E isomers, Z isomers, syn- isomers, anti- isomers, tautomers and the like. Aryl, heterocyclo or heteroaryl tautomers, heteroatom isomers and ortho, meta or para substitution isomers are also included as isomers. Solvates or solvent addition compounds such as hydrates or alcoholates are also specifically included both as chemicals of this invention and in, for example, formulations or pharmaceutical compositions for delivery.

Best Mode for Carrying Out the Invention


Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limiting of the remainder of the disclosure in any way whatsoever.

Abbreviations are often used for reagents and solvents in the specific examples that follow. Those abbreviations and their meanings are as follows:

- 5 BOC = t-butoxycarbonyl
 DEAD = diethyl azodicarboxylate
 DMF = dimethylformamide
 DMPU = 1,3-dimethyl-3,4,5,6-tetrahydro-
 2(1H)-pyrimidinone
10 EtOAc = ethyl acetate
 EDC = 1-ethyl-3-[3-(dimethylamino)-
 propyl]carbodiimide hydrochloride
 Et₂O = diethyl ether
 HOBT = 1-hydroxybenzotriazole
15 MeOH = methanol
 MeCl₂ = methylene chloride
 MsCl = methanesulfonyl chloride
 NMMOL or NMM = N-methyl morpholine
 THF = tetrahydrofuran
20 TsCl = toluenesulfonyl chloride
 THP-O-hydroxylamine = O-tetrahydropyran-
 hydroxylamine and O-tetrahydro-2H-
 pyran-2-yl-hydroxylamine
 Z = carbobenzyloxycarbonyl

25

Example 1: Preparation of N-hydroxy-3- [(4-methoxyphenyl)sulfonyl]-2-
 [(methylcarbonyl)amino]propanamide




Part A : A 250 mL round bottom flask equipped with magnetic stir bar and N₂ inlet was charged with 10.0 g N-acetyl dehydroalanine methyl ester, 9.9 g p-methoxythiophenol, and 10 mL NEt₃ in 100 mL of degassed MeOH. The reaction mixture was stirred one hour then concentrated *in vacuo* to remove NEt₃. The crude residue was taken up in 200 mL MeOH-150 mL H₂O and treated with 89 g (2.0 eq) Oxone®. The reaction was stirred 1 hour, filtered and the filtrate was concentrated *in vacuo*. Partitioning between EtOAc-water was followed by washing with water, brine, and concentration *in vacuo*. The crude off white solid was triturated with Et₂O to yield 14 g pure ester sulfone.

Part B: A 100 mL round bottom flask equipped with magnetic stir bar and N₂ inlet was charged with 500 mg ester sulfone, 1.2 mL (2.0 eq.) 50 % aqueous hydroxylamine in 8 mL MeOH/8 mL THF. The reaction was stirred overnight (about 18 hours) then concentrated *in vacuo* to a solid. Trituration with EtOAc yielded 380 mg of the desired hydroxamate.

Example 2: Preparation of 2-[[[(1,1-dimethylethyl)-amino]carbonyl]amino]-N-hydroxy-3-[(4-methoxyphenyl)sulfonyl]propanamide

-91-



Part A: A 250 mL round bottom flask equipped with magnetic stir bar and N₂ inlet was charged with
5 7.4 product from Example 1, Part A in 50 mL acetic acid and 50 mL concentrated HCl. The reaction was heated at reflux for 4 hours. The reaction mixture was concentrated *in vacuo* to a off white solid that was chased 2X with toluene, then vacuum dried to
10 yield 6.55 g of the amino acid HCl salt.

Part B : A 250 mL round bottom flask equipped with magnetic stir bar and N₂ inlet was charged with 6.5 g of the product amino acid HCl salt from above and 2.5 mL (1.5 eq.) of SOCl₂ in 70 mL MeOH. The
15 reaction was heated at reflux for 3 hours, then concentrated *in vacuo* to a crude solid. Trituration with EtO₂ yielded 6.9 g of amino ester HCl salt. The free base was generated using EtOAc-aqueous bicarbonate and was used without further
20 purification.

Part C: A 100 mL round bottom flask equipped with magnetic stir bar and N₂ inlet was charged with 900 mg of the amino ester from above and 288 mg t-butyl isocyanate in 20 mL MeCl₂. The reaction was
25 stirred overnight (about 18 hours), then concentrated *in vacuo*. Partitioning between hot EtOAc-water followed by washing with water, brine, and

SUBSTITUTE SHEET (RULE 26)


-92-

concentration *in vacuo* yielded 850 mg pure ester urea.

Part D: A 100 mL round bottom flask equipped with magnetic stir bar and N₂ inlet was charged with
5 90 mg ester urea from above and 0.2 mL (12.0 eq) 50 % aqueous hydroxylamine in 2 mL MeOH/2 mL THF. The reaction was stirred overnight (about 18 hours), then concentrated *in vacuo* to a solid. Trituration with Et₂O and filtration gave 50 mg of pure hydroxamate.

10

Example 3: Preparation of N-hydroxy-3-[(4-methoxyphenyl)sulfonyl]-2-
[(phenylcarbonyl)amino]propanamide



15

Part A : A 500 mL round bottom flask equipped with magnetic stir bar and N₂ inlet was charged with 5.0g (15.9 mmol) of ester from Example 1, Part A, in
20 50 mL acetic acid and 50 mL concentrated HCl. The reaction was heated at reflux for 4 hours. The reaction mixture was concentrated *in vacuo* to a white solid, then added and stripped with two aliquots of water and then two aliquots of toluene, then vacuum
25 dried to yield 4.6 g of pure amino acid HCl salt, m/z= 260 (M+H), as the HCl salt. m/z= 260.

Part B : A 500 mL round bottom flask equipped with magnetic stir bar and N₂ inlet was charged with

SUBSTITUTE SHEET (RULE 26)


4.6 g (15.7 mmol) amino acid HCl salt product from Part A in 100 mL MeOH. The solution was cooled in an ice bath, then treated with 1.7 mL (2.8 g, 23.5 mmol) thionyl chloride. The reaction was heated at reflux
5 for 3 hours, then concentrated in vacuo. The residue was partitioned between EtOAc and saturated sodium bicarbonate solution, the layers separated, and the aqueous layer was extracted with EtOAc (1X). The organic layers were combined and washed with brine,
10 dried (MgSO_4), and concentrated in vacuo to yield 2.9 grams of product suitable for the next reaction.

Part C : A 100 mL round bottom flask equipped with magnetic stir bar and N_2 inlet was charged with 0.7 g (2.7 mmol) product from Part B and 0.6 mL (0.4
15 g, 4.0 mmol) triethylamine in 10 mL CH_2Cl_2 . The solution was cooled in an ice bath and treated with 0.4 mL (0.5 g, 3.2 mmol) benzoyl chloride. After stirring for 1 hour, the reaction was concentrated in vacuo, the residue was partitioned EtOAc and water,
20 and the aqueous layer was extracted with EtOAc (1X). The organic layers were combined, washed with 5% citric acid solution, saturated bicarbonate solution and brine, dried (Na_2SO_4), and concentrated in vacuo to yield 0.8 g of crude amide product. The crude
25 amide product was chromatographed on silica gel using 50% ethyl acetate/hexane to yield 0.6 g of pure amide, $m/z = 378$ (M+H).

Part D: A 250 mL round bottom flask equipped with magnetic stir bar and N_2 inlet was charged with
30 0.6 g (1.6 mmol) of pure amide from Part C in THF (20 mL). The solution was treated with 0.5 mL (0.5 g, 8.21 mmol) 50 % aqueous hydroxylamine. After stirring for 65 hours, the reaction was not complete, and 0.1 g (4.3 mmol) of sodium metal and 5 mL MeOH

were added to the reaction to drive it to completion. The reaction was quenched with dry ice and then concentrated *in vacuo*. The residual solids were partitioned between EtOAc/2 N HCl solution. The solid precipitate was isolated by filtration, washed with water and EtOAc, then dried to yield 0.3 g of pure hydroxamate, $m/z = 379$ (M+H).

Example 4: Preparation of N-hydroxy-3-[(4-methoxyphenyl)sulfonyl]-2-
[(butylcarbonyl)amino]propanamide



Part A : A 500 mL round bottom flask equipped with magnetic stir bar and N₂ inlet was charged with 5.0 g (15.9 mmol) product from Example 1, Part A, in 50 mL acetic acid and 50 mL concentrated HCl. The reaction was heated at reflux for 4 hours. The reaction mixture was concentrated *in vacuo* to a white solid, and was then stripped with two aliquots of water and then two aliquots of toluene, then vacuum dried to yield 4.6 g of pure amino acid, $m/z = 260$ (M+H), as the HCl salt. $m/z = 260$.

Part B: A 250 mL round bottom flask equipped with magnetic stir bar and N₂ inlet was charged with 6.5 g product the amino acid from above and 2.5 mL (1.5 eq.) of SOCl₂ in 70 mL MeOH. The reaction was heated at reflux for 3 hours then concentrated in


vacuo to provide a crude solid. Trituration with EtO₂ yielded 6.9 g of amino ester HCl salt suitable for the next reaction.

Part C: A 100 mL round bottom flask equipped
5 with magnetic stir bar and N₂ inlet was charged with
1.0 g (3.2 mmol) of amino ester HCl salt product from
Part B and 1.4 mL (1.0 g, 9.7 mmol) triethylamine in
10 mL CH₂Cl₂. The reaction slurry was cooled in an
ice bath and treated with 0.5 mL (0.5 g, 3.9 mmol)
10 valeryl chloride. After stirring for 1 hour, the
reaction was incomplete, and an additional 0.5 mL
triethylamine and 0.4 mL valeryl chloride were added.
The reaction was stirred for 2 more hours, then
concentrated in vacuo. The residue was partitioned
15 EtOAc and water, and the aqueous layer was extracted
with EtOAc (1X). The organic layers were combined,
washed with 5% citric acid solution, saturated
bicarbonate solution and brine, dried (Na₂SO₄), and
concentrated in vacuo to yield 1.3 g of crude
20 product. The crude product was slurried with diethyl
ether, filtered and dried to yield 0.8 g of pure
amide, m/z = 358 (M+H).

Part D: A 100 mL round bottom flask equipped
with magnetic stir bar and N₂ inlet was charged with
25 0.8 g (2.3 mmol) of pure amide from Part C in THF (20
mL) and MeOH (10 mL). The solution was treated with
2.5 mL (1.4 g, 46.4 mmol) 50% aqueous hydroxylamine.
After stirring for 16 hours, the reaction mixture was
concentrated in vacuo, the residue was partitioned
30 between EtOAc/water, the layers were separated, and
the organic layer was washed with brine, dried
(Na₂SO₄), and concentrated in vacuo to yield a white
solid. The solid was slurried with diethyl ether,

isolated by filtration, then dried to yield 0.6 g of pure hydroxamate, $m/z = 359$ (M+H).

Example 5: Preparation of N-hydroxy-3-[(4-bromophenyl)sulfonyl]-2-[(methyl-carbonyl)amino]propanamide



Part A: A 1000 mL round bottom flask equipped with magnetic stir bar and N_2 inlet was charged with 25.0 g (132.2 mmol) 4-bromothiophenol and 19.9 g (138.8 mmol) methyl acetamidoacrylate in 400 mL degassed MeOH. The solution was cooled in an ice bath and treated with 19.4 mL (14.0 g, 138.8 mmol) triethylamine. The reaction was stirred for 1 hour and a white solid precipitated. The solid was isolated by filtration, washed with MeOH and dried to yield 27.6 g of product suitable for the next reaction.

Part B : A 1000 mL round bottom flask equipped with magnetic stir bar and N_2 inlet was charged with 11.0 g (33.1 mmol) of product from Part A in 400 mL MeOH. The slurry was warmed to yield a solution, then treated with 40 mL water and 61.1 g (99.3 mmol) Oxone®. The reaction mixture was stirred 4 hours, then filtered. The filter cake was washed with MeOH, and the filtrate was concentrated in vacuo. The residue was partitioned between EtOAc and H_2O , the


SUBSTITUTE SHEET (RULE 26)

- 97 -

layers were separated and the aqueous layer was extracted with EtOAc (2X). The organic phases were combined, washed with saturated sodium bicarbonate solution and brine, dried (MgSO₄), and concentrated in vacuo to yield the 6.7 g of pure product, m/z= 370 (M+Li).

Part C: A 100 mL round bottom flask equipped with magnetic stir bar and N₂ inlet was charged with 0.7 g (1.9 mmol) solid from Part B in THF (5 mL) and MeOH (5 mL). The solution was treated with 2.1 mL (1.2 g, 38.4 mmol) 50% aqueous hydroxylamine. After stirring for 17 hours, the reaction was concentrated in vacuo, the residue was partitioned between EtOAc/water, giving a three-phase system--the organic, aqueous and a solid precipitate at the interface. The solid was isolated by filtration, washed with diethyl ether, isolated by filtration, then dried to yield 0.2 g of pure hydroxamate, m/z= 387 (M+Na).

Example 6: Preparation of phenylmethyl-[2-[[2-(hydroxyamino)-2-oxo-1-[[[4-methoxyphenyl]sulfonyl]methyl]ethyl]amino]-2-oxoethyl]carbamate



Part A : A 100 mL round bottom flask equipped with magnetic stir bar and N₂ inlet was charged with


SUBSTITUTE SHEET (RULE 26)

-98-

1.0 g Z-Gly-OH, 920 mg (1.0 eq) EDC, 960 mg (1.5 eq) HOBT in 8 mL dry DMF. The reaction was activated for 10 minutes and then treated with 1.0 g free amino ester from Example 2, Part B. After stirring for 17 hours, the reaction was partitioned between EtOAc/saturated sodium bicarbonate. The combined organic phases were washed with 10 % KHSO₄, water, brine, and concentrated in vacuo to yield 1.25 grams of amide ester (>95 % by HPLC).

Part B : A 100 mL round bottom flask equipped with magnetic stir bar and N₂ inlet was charged with 700 mg of amide ester, 2.0 mL 50 % aq. hydroxylamine in 5 mL THF. The reaction was stirred 2 hours then quenched with dry ice. Standard work-up following evaporation from MeCl₂ gave the desired hydroxamate as a dry foam (>97 % by HPLC).

Example 7: Preparation of 4-[[1-[(hydroxyamino)-carbonyl]-2-[(4-methoxyphenyl)sulfonyl]-ethyl]amino-4-oxobutanoic acid
benzyl ester



Part A: A reaction mixture was prepared containing 2.36 g mono benzyl succinic acid (1.15 eq.), 2.18 g (1.15 eq.) EDC, and 2.0 g (1.5 eq.) HOBT in 8 mL dry of DMF. The reaction was activated for 10 minutes and then treated with 2.7 g free amino


SUBSTITUTE SHEET (RULE 26)

-99-

ester from Example 2, Part B. After stirring for 17 hours, the reaction was partitioned between EtOAc/saturated sodium bicarbonate. The combined organic phases were washed with 10 % KHSO₄, water, 5 brine, and concentrated *in vacuo* to yield 3.8 g of crude amide (82 % by HPLC). Flash chromatography with EtOAc yielded 3.1 g of pure diester product.

Part B: A 100 mL round bottom flask equipped with magnetic stir bar and N₂ inlet was charged with 10 436 mg of pure diester, and 1.3 mL 50 % aq. hydroxylamine in 20 mL MeOH and 5 mL THF. The reaction was stirred overnight (about 18 hours) then concentrated to provide a crude product (35 % by HPLC). Partitioning between EtOAc/water gave 200 mg 15 crude product (75 % by HPLC). Flash chromatography with 100 % THF gave 100 mg of pure hydroxamate.

Example 8: Preparation of 4[[1-[(hydroxyamino)-
carbonyl]-2-[(4-methoxyphenyl)sulfonyl]-
20 ethyl]amino-4-oxobutanoic acid



Part A: A 300 mL Paar flask equipped with 25 magnetic stir bar was charged with 436 mg of diester from Example 7, Part A, 100 mg 10 % Pd-C, in 35 mL MeOH, 30 mL AcOH then 60 mL THF. The reaction mixture was hydrogenated at 50 psi overnight (about 18 hours). After a N₂ purge the reaction was filtered


SUBSTITUTE SHEET (RULE 26)

-100-

through Celite® and concentrated to crude oil. The product was chased 2X with toluene then triturated with Et₂O to yield 630 mg (78 %) of the ester as a white solid.

5 Part B: A 100 mL round bottom flask equipped with magnetic stir bar and N₂ inlet was charged with 373 mg of the ester of Part A, and 1.0 mL 50 % aq. hydroxylamine in 4 mL MeOH and 5 mL THF. The reaction was stirred overnight (about 18 hours),
10 concentrated to a crude off white solid that was chased 2X with toluene then triturated with Et₂O to yield 250 mg of the title hydroxamate.

15 Example 9a: Preparation of O-(4-phenoxyphenyl)-
 N,N-dimethylcarbamothioate




 According to the general procedure described by
20 Newman [Org. Syn., 51, 139-142], 4-phenoxyphenol (150 mmol, 27.9 g) was added to an zero degrees C solution of potassium hydroxide (150 mmol, 8.4 g) in water (100 mL) under an argon atmosphere. After dissolution of the phenol, dimethylthiocarbamoyl
25 chloride (150 mmol, 24.8 g) in tetrahydrofuran (THF; 40 mL) was introduced to the reaction vessel at such a rate the internal temperature of the flask did not exceed 12 °C (over a period of about 20 minutes). After complete addition, the cooling bath was
30 removed, and stirring was continued an additional 10

SUBSTITUTE SHEET (RULE 26)

minutes. Aqueous KOH (10%, 50 mL) was added, and the reaction mixture was extracted with toluene (3 X 100mL). The combined organic layers were dried over magnesium sulfate, filtered through a short plug of silica, concentrated, seeded with a previous sample, and stored at zero degrees C. The resulting solid was triturated with methylene chloride (10 mL), affording 25.9 g of the title compound as a solid (63%). DSC 1 degree C/minute: 52.5-56.2 ° C.


10

Example 9b: Preparation of S-(4-phenoxyphenyl)-
N,N-dimethylcarbamothioate



15 The product from Example 9a (25.91 g) was heated under an argon atmosphere to 32 degrees C (internal) and maintained at that temperature for 15 minutes. The dark oil was allowed to cool, was characterized spectroscopically, and was used directly in the next
20 step.

Example 9c: Preparation of 4-(phenoxy)benzenethiol




25

Ethylene glycol (75 mL), water (10 mL) and potassium hydroxide (9 g) were added to the above-produced dark oil, and the mixture was heated at

reflux for 1 hour. Upon cooling, the mixture was poured over ice (about 200 mL), and washed with chloroform (2 X 150 mL). The aqueous layer was acidified with concentrated HCl, and the product was
5 extracted into chloroform (3 X 75 mL). The organic layers were collected, dried (MgSO₄), filtered through a short silica plug, and concentrated to afford the title compound as an oil (9.5 g, 49% from Example 9b)


- 10 Example 9d: Preparation of N-acetyl-3-[(4-phenoxyphenyl)sulphonyl]alanine, methyl ester



- 15 The product from Example 9c (15 mmol, 3.03 g) was diluted with methanol. Methyl 2-(acetamino)-2-propenoate (15 mmol, 2.14 g) was added, followed by triethylamine (15 mmol, 2.1 mL). The solution was stirred for one hour at ambient temperature, and for
20 2 hours at 45-5zero degrees C. The crude reaction mixture was rinsed into larger flask with a total of 50 mL additional methanol and cooled to zero degrees C. A solution of Oxone® (30 mmol, 18.45 g) in water (100 mL) was added. After 5 minutes, the cooling
25 bath was removed. The mixture was stirred at ambient temperature for one hour, then diluted with additional water (200 mL), and extracted with chloroform (200 mL, then 2 X 50 mL). The combined organic layers were dried (magnesium sulfate),


filtered, concentrated, and subjected to chromatography (hexane:ethyl acetate 1:1), affording the title compound as a white solid (2.185 g, 39%). DSC 1 degrees C/minute: 127.0-129.8. The product was
5 characterized spectroscopically. Elemental Anal. Calcd. for $C_{18}H_{19}NO_6S$: C, 57.28; H, 5.07; N, 3.71. Found: C, 57.10; H, 4.94; N, 3.51.

10 Example 9e: Preparation of N-acetyl-3-[(4-phenoxyphenyl)sulphonyl]alanine




The product from Example 9d (2.1 mmol, 800 mg)
15 was diluted with acetic acid (2 mL) and concentrated hydrochloric acid (2 mL) and warmed to 7zero degrees C for one hour. The reaction mixture was concentrated, and azeotroped with toluene (3 X 5 mL), affording the title compound as a solid (quant.). MS
20 MH^+ calcd. for $C_{17}H_{17}NO_6S$ 364, found 364. DSC 1 degree C/minute: 193.0-195.1. The product was characterized spectroscopically.

25 Example 9f: Preparation of N²-acetyl-3-[(4-phenoxyphenyl)sulphonyl]-N¹-(3,4,5,6-tetrahydro-2H-pyran-2-yl)alaninamide



The acid from Example 9e (2.1 mmol) was combined with hydroxybenzotriazole (2.7 mmol, 365 mg), O-tetrahydropyranhydroxylamine (3 mmol, 0.35 g), N,N-dimethylformamide (4 mL) and N-methylmorpholine (13 mmol, 1.43 mL). EDC (2.7 mmol, 518 mg) was added and the mixture was stirred at ambient temperature overnight (about 18 hours). The mixture was diluted with saturated NaHCO₃ (20 mL) and extracted with ethyl acetate (100 mL). The organic phase was washed with brine (20 mL), then dried over magnesium sulfate, filtered, concentrated, and subjected to silica gel chromatography (6% methanol in chloroform), affording the title compound as a glass (472 mg, 50%). The product was characterized spectroscopically.

Example 9g: Preparation of N²-acetyl-N¹-hydroxyl-3-
[(4-phenoxyphenyl)sulphonyl]alaninamide




The product from Example 9f (1.0 mmol, 450 mg) was dissolved in dioxane (5 mL). Six normal HCl (0.5

mL) was added, and the mixture was stirred one hour. Concentration followed by chromatography afforded the title compound as a glass (120 mg, 32%). MS MH⁺ calcd. for C₁₇H₁₈N₂O₆S 378, found 379

5


Example 10a: Preparation of 3-[(4-phenoxyphenyl)-sulphonyl]alanine, monhydrochloride



10 A solution of the titled compound of Example 9d (3.27 g, 8.66 mmol) in glacial acetic acid (18.5 mL) was heated until all solids dissolved and then concentrated HCl (18.5 mL) was added. The solution was heated at reflux for 4 hours. The mixture was
15 concentrated to an off white solid. The solid was treated with toluene, concentrated, and triturated with diethyl ether afforded the title amino acid as a white solid (3.04 g, 98%). Elemental Anal. Calcd. for C₁₅H₁₆NO₅SCl: C, 50.35; H, 4.51; N, 3.91; S, 8.96.
20 Found: C, 50.21; H, 4.43; N, 3.92; S, 9.04.


Example 10b: Preparation of N-[(1,1-dimethylethyl)-carbonyl]-3-[(4-phenoxyphenyl)-alaninamide]

25




The amino acid (1.4 mmol, 500 mg) from above was dissolved in a mixture of water (2 mL), triethylamine (5.6 mmol, 0.79 mL), and acetone (3 mL), and cooled to zero degrees C. Pivaloyl chloride (1.8 mmol, 0.22 mL) was added over 5 minutes. The reaction was warmed to ambient over one hour and stirred overnight (about 18 hours), then concentrated and azeotroped with acetonitrile (10 mL). Chromatography (20% methanol/chloroform) afforded the title compound as the triethylamine salt. The triethylamine salt was dissolved in chloroform (150 mL) and washed with 2N HCL (20 mL). The organic layer was dried over magnesium sulfate, filtered, concentrated and re-chromatographed to afford the title compound as a white foam (485 mg). The product was characterized spectroscopically.

Example 10c: Preparation of N²-(1,1-dimethylethyl)-3-[(4-phenoxyphenyl)sulphonyl]-N¹-3,4,5,6-tetrahydro-2H-pyran-2-yl)alaninamide



The acid from Example 10b (0.89 mmol, 455 mg) was suspended in tetrahydrofuran. EDC (1.00 mmol, 192 mg) was added. After 15 minutes, THP-O-hydroxylamine was added. The mixture was stirred 20 hours at ambient temperature, diluted with saturated sodium bicarbonate, and extracted with ethyl acetate (125 mL). The organic layer was washed with brine (5 mL), dried over magnesium sulfate, filtered, concentrated and subjected to chromatography (6% methanol/chloroform) to afford the title compound as a solid (260 mg, 58%). The product was characterized spectroscopically. Elemental Anal. Calcd. for $C_{24}H_{31}N_2O_7S(1/4 H_2O)$: C, 59.10; H, 6.25; N, 5.51. Found: C, 59.10; H, 6.50; N, 5.31.


Example 10d: Preparation of N²-[(1,1-dimethylethyl)-carbamoyl]-N¹-hydroxy-3-[(4-phenoxyphenyl)-sulphonyl]alaninamide



The product of Example 10c (, 250 mg) was suspended in methanol (10 mL) and toluenesulphonic acid (10 mg) was added. After 2 hours, additional toluenesulphonic acid (10 mg) was added, and stirring was continued an additional hour. The reaction mixture was concentrated, and the residue was subjected to chromatography (6% methanol/chloroform), affording the desired hydroxamate as a white foam (166 mg). The product was characterized spectroscopically. Elemental Anal. Calcd. for $C_{20}H_{24}N_2O_6S$: C, 57.13; H, 5.75; N, 6.66. Found: C, 52.67; H, 5.32; N, 5.91.

15

Example 11a: Preparation of N-acetyl-3-[(4-fluorophenyl)sulfonyl]alanine, methyl ester




20

To a solution of methyl 2-acetamidoacrylate (10.0 g, 69.9 mmol) in 100 mL of methanol, cooled to

zero degrees C, was added 4-fluorothiophenol (7.44 mL, 69.9 mmol) followed by triethylamine (9.74 mL, 69.9 mmol). The mixture stirred under a nitrogen atmosphere for 4 hours. The solution was poured into
5 a slurry of Oxone[®] (86 g, 140 mmol) in H₂O (250 mL) and MeOH (125 mL). After stirring at room temperature for 18 hours, the mixture was partitioned between EtOAc (200 mL) and H₂O and washed with saturated NaHCO₃ (3 x 100 mL) and brine (1 x 100 mL).
10 After concentration *in vacuo*, trituration with diethyl ether afforded the title compound as a white solid (16.3 g, 77%): MS MH⁺ calcd. for C₁₂H₁₄NO₅SF: 304, found 304. HRMS calcd. for C₁₂H₁₄NO₅SF: 303.0577, found: 303.0581.

15


Example 11b: Preparation of N-acetyl-3-[(4-fluorophenyl)sulfonyl]alanine



20

A solution of the title compound of Example 11a (17.9 g, 59.0 mmol) in glacial acetic acid (135 mL) and concentrated HCl (135 mL), was heated to 7 degrees C for 2 hours. The solution was concentrated
25 under a stream of N₂ to yield the title compound as a white solid (17.0 g, quant. yield): MS MH⁺ calcd. for C₁₁H₁₂NO₅SF: 290, found 290. HRMS calc. for C₁₁H₁₃NO₅SF: 290.0499, found: 290.0520.

Example 11c: Preparation of N-acetyl-3-[[4-(phenylthio)phenyl]sulfonyl]alanine




5

Thiophenol (7.27 g, 71.0 mmol) was added to a solution of the title compound of Example 11b (17.0g, 59.0 mmol) and K₂CO₃ (24.4 g, 177 mmol) in DMF (200 mL). After heating at 9 degrees C for 2 hours, the mixture was concentrated under a stream of N₂. The solution was acidified to pH=1 with 1M KHSO₄, extracted into EtOAc (2X) and the organic phases were washed with brine and dried over MgSO₄. The solution was concentrated *in vacuo* to yield the title compound as a white solid and was carried on without additional purification.

15

Example 11d: Preparation of 3-[[4-(phenylthio)-phenyl]sulfonyl]alanine, monohydrochloride

20



A solution of the title compound of Example 11c (30.0 g) in glacial acetic acid (100 mL) and concentrated HCl (100 mL) was heated to 11 degrees C for 19 hours. The solution was concentrated and


25

SUBSTITUTE SHEET (RULE 26)

-111-

trituration with diethyl ether yielded the title compound as a white solid (17.1 g, 78% for 3 steps):
Anal. Calcd. for $C_{15}H_{15}NO_4S_2 \cdot 1.1 HCl \cdot 0.25 H_2O$: C, 47.16;
H, 4.38; N, 3.67; Cl, 10.21. Found: C, 47.16; H,
5 4.27; N, 3.91; Cl, 9.95.

Example 11e: Preparation of N-(1-oxopentyl)-3-[[4-(phenylthio)phenyl]sulfonyl]alanine




10

To a solution of the title compound of Example 11d (1.0 g, 2.68 mmol) in H_2O (3 mL) and acetone (2 mL), cooled to zero degrees C, was added
15 triethylamine (1.17 mL, 8.43 mmol), followed by drop-wise addition of valeryl chloride (0.30 mL, 2.52 mmol) in acetone (2 mL). After stirring at room temperature for 18 hours, the mixture was concentrated. The resulting aqueous residue was
20 acidified with 1M $KHSO_4$ and extracted with CH_2Cl_2 (3X). The combined organic phases were washed with 1M $KHSO_4$ (1 x 25 mL) and brine (1 x 25 mL) and dried over $MgSO_4$. The solution was concentrated in vacuo to yield the title compound as a colorless oil (690 mg,
25 61%): MS M^+ calcd. for $C_{20}H_{23}NO_5S_2$: 421, found 421. HRMS calc for $C_{20}H_{23}NO_5S_2$: 421.1018, found: 421.1025.

Example 11f: Preparation of N-[2-oxo-1-[[[4-(phenylthio)phenyl]sulfonyl]methyl]-2-

SUBSTITUTE SHEET (RULE 26)

[[[(3,4,5,6- tetrahydro-2H-pyran-2-yl) -
oxy] amino] ethyl] pentanamide




5

To a solution of the title compound of Example 11e (690 mg, 1.64 mmol) and O-tetrahydro-2H-pyran-2-yl-hydroxylamine (192 mg, 1.64 mmol) in anhydrous THF (40 mL), cooled to zero degrees C, was added EDC (330 mg, 1.72 mmol) and HOBT (264 mg, 1.72 mmol). After stirring at zero degrees C for 5 hours and at room temperature for 18 hours, the solution was concentrated. The residue was dissolved into EtOAc (50 mL) and washed with 1M KHSO₄ (2 x 50 mL), saturated NaHCO₃ (2 x 50 mL), and brine (1x 50 mL) and dried over Na₂SO₄. Purification by chromatography (1:1 EtOAc/hexane) yielded the title compound as a colorless oil (500 mg, 59%): Anal. Calcd. for C₂₅H₃₂N₂O₆S₂: C, 57.67; H, 6.19; N, 5.38. Found: C, 57.48; H, 6.37; N, 5.11.

20


Example 11g: Preparation of N-[1-[(hydroxyamino)-carbonyl]-2-[[4-(phenylthio)phenyl]-sulfonyl]ethyl]pentanamide

25



To a solution of the title compound of Example 11f (500 mg, 0.96 mmol) in methanol (20 mL) was added p-toluenesulfonic acid, monohydrate (40 mg, 0.21 mmol). After stirring at room temperature for 2 hours, the solution was concentrated. Purification by chromatography (6:94 MeOH/CHCl₃) yielded the title compound as a light beige solid (170 mg, 40%): Anal. Calcd. for C₂₀H₂₄N₂O₅S₂·0.5 H₂O: C, 53.91; H, 5.66; N, 6.29; S, 14.39. Found: C, 53.98; H, 5.44; N, 6.20; S, 14.47.

Example 12a: Preparation of




15

A solution of p-toluenesulfonyl chloride (TsCl; 0.779 g, 4.19 mmol) in 2.4 mL acetone was added dropwise to a zero degree C mixture of the title compound of Example 12a (1.50 g, 4.19 mmol) and triethylamine (2.36 mL, 16.9 mmol) in 2.4 mL acetone and 6.4 mL H₂O. After 3 hours at room temperature, the mixture was concentrated, diluted with toluene, and concentrated to give a white solid. The solid was washed with water and toluene, and then purified by

chromatography (5:95 MeOH/CHCl₃) to afford the depicted compound as a white solid (0.75 g, 38 % yield). ¹H NMR spectrum was consistent for the depicted compound.


5

Example 12b: Preparation of



10 To a solution of the depicted compound of Example 12a (0.75 g, 1.6 mmol), HOBT (0.253 g, 1.7 mmol), and O-tetrahydro-2H-pyran-2-yl-hydroxylamine (0.18 g, 1.6 mmol) in dry THF (22 mL) at zero degrees C was added EDC (0.316 g, 1.7 mmol). The mixture was
15 slowly warmed to room temperature and stirred overnight (about 18 hours) at room temperature. The solution was concentrated, diluted with ethyl acetate, and washed with H₂O and brine. After drying over MgSO₄, the filtrate was concentrated to a white
20 solid. The solid was purified by chromatography (40:60 acetone/hexane) to give the title compound as a white solid (0.22 g, 24 % yield). The proton NMR spectrum was consistent for the title compound.


25 Example 12c: Preparation of



SUBSTITUTE SHEET (RULE 26)

A solution of the depicted compound of Example 12b (0.220 g, 0.383 mmol) and *p*-toluenesulfonic acid monohydrate (20 mg, 0.105 mmol) in MeOH (5 mL) was stirred at room temperature for 2 hours. The solution was concentrated and triturated with ET₂O to afford the depicted compound as an off white solid (50 mg, 43 %): MS M⁺ calcd. for C₂₂H₂₂N₂O₇S₂ 490, found 490. HRMS calcd. for C₂₂H₂₂N₂O₇S₂: 490.0868; found: 490.0863.


Example 13a: Preparation of 3-[(4-phenoxyphenyl)-sufonyl]-N-[(phenylmethoxy)-carbonyl]alanine



The title compound was prepared by the method of Example 10b, using (0.892 g, 2.5 mmol) of the amino acid of 34a, and using carbobenzyloxy chloride (0.43 mL, 3 mmol) in the place of pivaloyl chloride. The product was a yellow foam (689 mg, 66%), and was characterized spectroscopically. MS MH⁺ calcd. for C₂₃H₂₁NO₇S 456, found 456. Elemental Anal. Calcd for C₂₃H₂₁NO₇S: C, 59.47; H, 4.77; N, 3.02. Found: C, 59.49; H, 4.35; N, 3.09.

Example 13b: Preparation of phenylmethyl[2-oxo-1-[[[(4-phenoxyphenyl)sulfonyl]methyl]-2-[[3,4,5,6-tetrahydro-2H-pyran-2-yl]-


oxy]amino]ethyl carbamate



5 The title O-THP hydroxamate was prepared from the acid from Example 15a (669 mg, 1.6 mmol) by the method of Example 10C, affording after chromatography 208 mg product as a yellow foam. The structure was confirmed spectroscopically.

10

Example 13c: Preparation of phenylmethyl [1-
[(hydroxyamino) carbonyl] -2- [(4-
phenoxyphenyl) sulfonyl] ethyl] carbamate



15

The O-THP hydroxamate from Example 13b (203 mg, 0.36 mmol) was diluted with methanol (10 mL), and toluenesulphonic acid (20 mg) was added. The mixture
20 was stirred 1 hour, concentrated, and subjected to chromatography, affording the title hydroxamate (74 mg, 20%) as a white foam, which was characterized


SUBSTITUTE SHEET (RULE 26)

-117-

spectroscopically. MS MH⁺ calcd. for C₂₃H₂₂N₂O₇S 470, found 470. Elemental Anal. Calcd for C₂₃H₂₂N₂O₇S (H₂O): C, 56.53; H, 4.90; N, 5.65. Found: C, 56.55; H, 4.95; N, 5.73.

5

Example 14a: Preparation of N-[(phenylmethoxy)-carbonyl]-3-[[4-(phenylthio)phenyl]-sulfonyl]alanine




10

To a solution of the title compound of Example 11d (1.5 g, 4.0 mmol) in H₂O (5 mL) and acetone (3 mL), cooled to zero degrees C, was added triethylamine (1.8 mL, 12.6 mmol), followed by dropwise addition of benzyl chloroformate (0.57 mL, 4.0 mmol) in acetone (3 mL). After stirring at room temperature for 6 hours, the mixture was concentrated. The resulting aqueous residue was acidified with 1M KHSO₄ and extracted with CH₂Cl₂ (3X). The combined organic phases were washed with 1M KHSO₄ (1 x 25 mL) and brine (1 x 25 mL), and dried over MgSO₄. The solution was concentrated *in vacuo* to yield the title compound as a yellow oil (1.4 g, 72%).

Example 14b: Preparation of phenylmethyl-[2-oxo-1-[[[4-(phenylthio)phenyl]sulfonyl]-

SUBSTITUTE SHEET (RULE 26)


methyl]-2-[[[3,4,5,6-tetrahydro-2H-pyran-2-yl]-oxy]amino]ethyl]carbamate



5


To a solution of the title compound of Example 14a (1.35 g, 2.87 mmol) and O-tetrahydro-2H-pyran-2-yl-hydroxylamine (336 mg, 2.87 mmol) in anhydrous DMF (20 mL), cooled to zero degrees C, was added EDC (578 mg, 3.01 mmol) and HOBT (461 mg, 3.01 mmol). After stirring at zero degrees C for 1 hour and at room temperature for 2 hours, the solution was partitioned between EtOAc and H₂O. The resulting organic phase was washed with 1M KHSO₄ (2 x 50 mL), saturated NaHCO₃ (2 x 50 mL) and brine (1x 50 mL), and dried over Na₂SO₄. Purification by chromatography (1:1 EtOAc/hexane) yielded the title compound as a colorless oil (500 mg, 31%):

20 Example 14c: Preparation of phenylmethyl-[2-(hydroxyamino)-2-oxo-1-[[[4-(phenylthio)-phenyl]sulfonyl]methyl]-ethyl]carbamate



To a solution of the title compound of Example 14b (500 mg, 0.96 mmol) in acetonitrile (20 mL) was added p-toluenesulfonic acid, monohydrate (40 mg, 0.21 mmol). After stirring at room temperature for 1.5 hours, the solution was concentrated. Trituration with diethyl ether yielded the title compound as a white solid (120 mg, 28%): Anal. Calcd. for $C_{23}H_{22}N_2O_6S_2 \cdot 0.2H_2O$: C, 56.36; H, 4.61; N, 5.72; S, 13.08. Found: C, 56.18; H, 4.21; N, 5.50; S, 13.31.

Example 15a: Preparation of 2-(acetylamino)-3-[[4-(phenylthio)phenyl]sulfonyl]-N-[(3,4,5,6-tetrahydro-2H-pyran-2-yl)oxy]propanamide




To a solution of the title compound of Example 11c (930 mg, 2.45 mmol) and O-tetrahydro-2H-pyran-2-yl-hydroxylamine (287 mg, 2.45 mmol) in anhydrous DMF (40 mL), cooled to zero degrees C, were added EDC

SUBSTITUTE SHEET (RULE 26)

(493 mg, 2.57 mmol) and HOBT (393 mg, 2.57 mmol). After stirring at zero degrees C for 1 hour and at room temperature for 3 hours, the solution was partitioned between EtOAc and H₂O. The resulting organic phase was washed with 1M KHSO₄ (2 x 50 mL), saturated NaHCO₃ (2 x 50 mL) and brine (1x 50 mL), and dried over Na₂SO₄. Purification by chromatography (1:1 EtOAc/hexane) yielded the title compound as a colorless oil (400 mg, 34%).

10

Example 15b: Preparation of 2-(acetylamino)-N-hydroxy-3-[[4-(phenylthio)phenyl]sulfonyl]propanamide




15

To a solution of the title compound of Example 15a (400 mg, 0.84 mmol) in methanol (40 mg), was added p-toluenesulfonic acid, monohydrate (40 mg, 0.21 mmol). After stirring at room temperature for 2 hours, the solution was concentrated. Purification by flash chromatography (4:96 MeOH/CHCl₃) yielded the title compound as a white solid (140 mg, 42%): Anal. Calcd. for C₁₇H₁₈N₂O₅S₂·0.8H₂O: C, 49.94; H, 4.83; N, 6.85; S, 15.68. Found: C, 50.23; H, 4.62; N, 6.80; S, 15.29.


25

Example 16a: Preparation of 3-[[4-(phenylthio)phenyl]sulfonyl]alanine, methyl ester



Thionyl chloride (0.44 mL, 6.0 mmol) was added
5 drop-wise to a solution of the title compound of
Example 11d (1.5 g, 4.0 mmol) in methanol (20 mL).
The solution was heated at reflux for 5 hours and
then concentrated. After adding EtOAc, the insoluble
material was removed by vacuum filtration. The
10 filtrate was washed with saturated NaHCO₃ (2 x 50 mL)
and brine (1 x 50 mL), and dried over MgSO₄. The
solution was concentrated to yield the title compound
as a yellow oil (570 mg, 41%):

15 Example 16b: Preparation of N-[2-[(phenylmethoxy)-
carbonyl]amino]acetyl]-3-[[4-
(phenylthio)phenyl]sulfonyl]alanine,
methyl ester




20

Carboxybenzyloxycysteine succinimidyl ester (500
mg, 1.6 mmol) was added to a solution of the title
compound of Example 16a (570 mg, 1.6 mmol) in
25 methylene chloride (20 mL) cooled to zero degrees C.

SUBSTITUTE SHEET (RULE 26)


After stirring at room temperature for 20 hours, the solvent was removed in vacuo, and the residue was dissolved into EtOAc. The organic layer was washed with 1M KHSO₄ (2 x 50 mL), saturated NaHCO₃ (2 x 50 mL), and brine (1 x 50 mL), and dried over MgSO₄. Concentration in vacuo yielded the title compound as a colorless oil (800 mg, 93%).

Example 16c: Preparation of phenylmethyl-[2-[[2-(hydroxyamino)-2-oxo-1-[[[4-(phenylthio)-phenyl]sulfonyl]methyl]-ethyl]amino]-2-oxoethyl]carbamate




15 To a solution of the title compound of Example
16b (440 mg, 0.81 mmol) in tetrahydrofuran (9 mL) and
methanol (3 mL) was added NH_2OH (50% aqueous solution,
0.74 mL, 12 mmol). After stirring for 20 hours at
20 room temperature, the solution was concentrated.
Purification by flash chromatography (5:95 MeOH/ CHCl_3)
yielded the title compound as a white solid (80mg,
18%): Anal. Calcd. for $\text{C}_{25}\text{H}_{25}\text{N}_3\text{O}_7\text{S}_2 \cdot 0.25 \text{ H}_2\text{O}$: C, 54.78;
H, 4.69; N, 7.67; S, 11.70. Found: C, 54.78; H,
25 4.72; N, 7.46; S, 11.53.

Example 17a: Preparation of



5 To a solution of the title compound of Example 11d (1.6 g, 4.28 mmol) in H₂O (6 mL) and acetone (3 mL) at zero degrees C was added triethylamine (2.8 mL, 38.2 mmol), followed by drop-wise addition of tosyl chloride (TsCl; 1.24 g, 65.0 mmol) in acetone
10 (3 mL). After stirring at room temperature for 18 hours, the mixture was concentrated. The residue treated with a 10% KHSO₄ aqueous solution and CH₂Cl₂ provided a suspension that was filtered to afford the desired as a white powder (1.3 g, 61.9%). HRMS
15 Calcd. For C₂₂H₂₁NO₆S₃: 492.0609. Found: 492.0607.

Example 17b: Preparation of




20

To a solution of the depicted compound of Example 17a (0.6 g, 1.22 mmol) in anhydrous THF (12 mL) and anhydrous DMF (3 mL) at zero degrees C were added EDC (0.280 g, 1.46 mmol) and HOBT (0.198 g, 1.46 mmol). After stirring at zero degrees C for 30
25 minutes, O-tetrahydro-2H-pyran-2-yl-hydroxylamine (0.172 g, 1.46 mmol) was added and the reaction


solution was stirred at room temperature for 16 hours. Then the solution was concentrated and the residue was dissolved into EtOAc (50 mL) and washed successively with 10% citric acid (2 x 50 mL),
5 saturated NaHCO_3 (2 x 50 mL) and brine (1x 50 mL), and dried over MgSO_4 . Purification by chromatography (1.5/60/40 MeOH/EtOAc/hexane) yielded the depicted compound as a white solid (0.47 g, 65.3%).

10 Example 17c: Preparation of



p-Toluenesulfonic acid monohydrate (47 mg, 0.25
15 mmol) was added to a solution of the depicted compound of Example 17b (430 mg, 0.72 mmol) in methanol (13 mL). After stirring at room temperature for 1.5 hours, a white precipitate appeared that was filtered to afford a white solid. Trituration with
20 ether gave the depicted compound (280 mg, 75.7%):
Anal. Calcd. for $\text{C}_{22}\text{H}_{22}\text{N}_2\text{O}_6\text{S}_3$: C, 51.12; H, 4.58; N, 7.95; S, 18.20. Found: C, 51.06; H, 4.59; N, 7.83; S, 18.45.


25 Example 18a: Preparation of



SUBSTITUTE SHEET (RULE 26)

To the title compound of 16a (0.5 g, 1.4 mmol) in MeOH (2 mL) and THF (2 mL) was added 50% aqueous NH_2OH (1 mL, 13.5 mmol). The reaction solution was stirred for 2 hours during which time a white precipitate appeared. The resulting suspension was filtered, and the white powder thus obtained was washed with ether (10 mL X 3) to afford the depicted compound. Anal. Calcd. For $\text{C}_{15}\text{H}_{16}\text{N}_2\text{O}_4\text{S}_2$: C, 51.12; H, 4.58; N, 7.95; S, 18.20. Found: C, 51.06; H, 4.59; N, 7.83; S, 18.45.

Example 19a: Preparation of




15

To a solution of Cbz-alanine (1.43 g, 6.4 mmol) in DMF at zero degrees C were added EDC (1.23 g, 6.4 mmol) and HOBT (0.87 g, 6.4 mmol). After 30 minutes at zero degrees C, the title compound of Example 11d (2g, 5.36 mmol) was added to the solution followed by N-methylmorpholine (1.82 mL, 16.5 mmol). The reaction solution was left stirring overnight (about 18 hours) at room temperature. The reaction was concentrated and the resulting residue was dissolved in aqueous 10% KHSO_4 solution. The aqueous solution was extracted with ethyl acetate and the combined

25


organic phases were washed with 10% KHSO₄ aqueous solution, dried over MgSO₄ and evaporated to afford the depicted compound.

5 Example 19b: Preparation of



To a solution of the depicted compound of
10 Example 19a (1.0 g, 1.84 mmol) in dry DMF (10 mL) at
zero degrees C were added EDC (0.422 g, 2.2 mmol) and
HOBT (0.299 g, 2.2 mmol). After stirring at zero
degrees C for 30 minutes, O-tetrahydro-2H-pyran-2-yl-
hydroxylamine (0.323 g, 2.6 mmol) was added. The
15 reaction solution was then stirred at room
temperature for 16 hours. Concentration of the
solution afforded a residue that was dissolved into
EtOAc (50 mL) and washed with 10% citric acid (2 x 50
mL), saturated NaHCO₃ (2 x 50 mL), and brine (1x 50
20 mL), and then dried over MgSO₄. Evaporation gave a
residue that was purified by chromatography on silica
gel (2/50/50 MeOH/EtOAc/hex) to afford the depicted
compound as white solid (0.6 g, 50.8%). Anal. calcd.
for C₃₁H₃₅N₃O₈S₂: C, 58.02; H, 5.5; N, 6.55; S, 9.99.
25 Found: C, 57.69; H, 5.59; N, 6.35; S, 10.33.


Example 19c: Preparation of



p-Toluenesulfonic acid monohydrate (25 mg, 0.13
5 mmol) was added to a solution of the title compound
of Example 19b (170 mg, 0.26 mmol) in methanol (1 mL)
and acetonitrile (4 mL). After stirring at room
temperature overnight (about 18 hours), the reaction
solution was concentrated to a residue that was
10 purified by chromatography on silica gel (5/100
MeOH/CH₂Cl₂) to afford the title compound as white
solid (82 mg, 55.5%): Anal. Calcd. for C₂₆H₂₇N₃O₇S₂: C,
56.00; H, 4.88; N, 7.54; S, 11.50. Found: C, 56.11;
H, 5.22; N, 7.28; S, 11.22.


15

Example 20a: Preparation of



The title compound of Example 16a (0.43 g, 1.2 mmol) in CH_2Cl_2 was added t-butyl isocyanate (0.48 mL, 4.2 mmol). After stirring for 3 days at room temperature, the reaction solution was diluted with ethyl acetate (50 mL) and washed successively with 10% citric acid (2 x 50 mL), saturated NaHCO_3 (2 x 50 mL), and brine (1x 50 mL) and then dried over MgSO_4 . Evaporation of the solvent gave the title compound as crystalline white solid. Anal. Calcd. For $\text{C}_{21}\text{H}_{26}\text{N}_2\text{O}_5\text{S}_2$:
C, 55.98; H, 5.82; N, 6.22; S, 14.25. Found: C, 55.78; H, 6.00; N, 6.09; S, 14.25.

Example 20b: Preparation of



15

Aqueous NH_2OH (0.9 mL, 15.8 mmol) was added to a solution of the depicted compound of Example 20a (0.47 g, 1.04 mmol) in MeOH (2 mL) and THF (2 mL). After stirring overnight (about 18 hours) at room temperature, the reaction solution was concentrated to a residue that was purified by chromatography on silica gel (5/100 MeOH/ CH_2Cl_2) to afford the depicted compound as a white solid (287 mg, 61.0%): Anal. Calcd. For $\text{C}_{20}\text{H}_{25}\text{N}_3\text{O}_5\text{S}_2 \cdot \text{H}_2\text{O}$: C, 51.16; H, 5.80; N, 8.95; S, 13.66. Found: C, 51.34; H, 5.60; N, 9.41; S, 13.57.

Example Compounds 21-70 are similarly prepared to those above.

5 Example 21: In Vitro Metalloprotease Inhibition

The compounds prepared in the manner described in the Examples above were assayed for activity by an *in vitro* assay. Following the procedures of Knight et al., *FEBS Lett.* 296(3):263
10 (1992). Briefly, 4-aminophenylmercuric acetate (APMA) or trypsin-activated MMPs were incubated with various concentrations of the inhibitor compound at room temperature for 5 minutes.

More specifically, recombinant human MMP-13
15 and MMP-1 enzymes were prepared in laboratories of the assignee following usual laboratory procedures. MMP-13 from a full length cDNA clone was expressed as a proenzyme using a baculovirus as discussed in V.A. Luckow, *Insect Cell Expression Technology*, pages 183-
20 218, in Protein Engineering: Principles and Practice, J.L.Cleland et al eds., Wiley-Liss, Inc., (1996). See, also, Luckow et al., *J. Virol.*, 67:4566-4579 (1993); O'Reilly et al., Baculovirus Expression Vectors: A Laboratory Manual, W.H. Freeman and
25 Company, New York, (1992); and King et al., The Baculovirus Expression System: A Laboratory Guide, Chapman & Hall, London (1992) for further details on use of baculovirus expression systems. The expressed enzyme was purified first over a heparin agarose
30 column and then over a chelating zinc chloride column. The proenzyme was activated by APMA for use in the assay.

MMP-1 expressed in transfected HT-1080 cells was provided by Dr. Harold Welgus of Washington

University, St. Louis, MO. The enzyme was also activated using APMA and was then purified over a hydroxamic acid column. Further specifics for preparation and use of these enzymes can be found in the scientific literature describing these enzymes. See, for example, Enzyme Nomenclature, Academic Press, San Diego, Ca (1992) and the citations therein, and Frije et al., J. Biol. Chem., 269(24): 16766-16773 (1994).

The enzyme substrate is a methoxycoumarin-containing polypeptide having the following sequence: MCA-ProLeuGlyLeuDpaAlaArgNH₂, wherein MCA is methoxycoumarin and Dpa is 3-(2,4-dinitrophenyl)-L-2,3-diaminopropionyl alanine. This substrate is commercially available from Baychem as product M-1895.

The buffer used for assays contained 100 mmol Tris-HCl, 100 mmol NaCl, 10 mmol CaCl₂ and 0.05 percent polyethyleneglycol (23) lauryl ether at a pH value of 7.5. Assays were carried out at room temperature, and dimethyl sulfoxide (DMSO) at a final concentration of 1 percent was used to dissolve inhibitor compound.

The assayed inhibitor compound in DMSO/buffer solution was compared to an equal amount of DMSO/buffer with no inhibitor as control using Microfluor™ White Plates (Dynatech). The inhibitor or control solution was maintained in the plate for 10 minutes and the substrate was added to provide a final concentration of 4 μM.

In the absence of inhibitor activity, a fluorogenic peptide was cleaved at the gly-leu peptide bond, separating the highly fluorogenic peptide from a 2,4-dinitrophenyl quencher, resulting

in an increase of fluorescence intensity (excitation at 328 nm/emission at 415 nm). Inhibition was measured as a reduction in fluorescent intensity as a function of inhibitor concentration, using a Perkin Elmer L550 plate reader. The IC₅₀ values were calculated from those values. The results are set forth in the Inhibition Table below, reported in terms of IC₅₀ values in nanomolar (nM) amounts.

10

Inhibition Table

MMP ENZYME INHIBITION PROFILE
(nM, IC₅₀)

15

Example	MMP-1	MMP-2	MMP-13
1	300		15.0
2	1500		15.0
3	2000		25.0
4	400		3.5
5	100		200
6	40		<1.0
7	200		2.5
8	600		60.0
9g	770		1.1
10d	1600		2.7
11g	10000		6.0
12c	400	<0.1	0.6
13c	1400		1.1
14c	>10000		3.7
15b	>10000	2.2	5.9
16c	>10000		0.8
17c	600		0.4

SUBSTITUTE SHEET (RULE 26)

18a	>10000		24.0
19a	>10000		0.8
20b	>10000		4.0
21	300		15.0
22	250		15.0
23	>10000		1500
24	5200		650
25	1.0		1.0
26			
27	>10000		14.0
28	>10000	5.5	13.0
29	>10000		7.0
30	10000	0.5	2.9
31	3000	0.8	0.4
32	8000		0.4
33	500	0.2	0.3
34	>10000		5.0
35	3000		0.2
36	>10000		2.5
37	350	0.25	0.4
38	170	0.5	0.5
39	114, 225	0.3, 0.8	0.3, 1.1
40	1600	0.3	0.3
41	440, 670	0.3, 0.5	0.4, 1.0
42	>10000	40.0	41.5
43	90.0	0.2	0.2
44	900	0.3	0.8
45	>10000		5.0
46	1310	0.3	1.1
47	258	0.6	45.3
48	475	<0.1	0.2
49	>10000	80.0	160
50	316	0.3	0.3

SUBSTITUTE SHEET (RULE 26)

-133-

51	295	0.3	0.4
52	>10000	5.0	24.0
53	8100	0.3	2.1
54	345	0.2	0.3
55	1600	1.3	1.6
56	1200	0.2	0.3
57	>10000	540	580
58	4400	3.2	2.4
59	580	0.4	0.8
60	440	0.3	0.4
61	>10000	1.1	60.0
62	72.3	23.9	1200
63	3050	1.0	21.5
64	7000	0.4	8.1
65	>10000	184	161
66	2350	0.5	1.1
67	2400	1.3	2.4
68	>10000	0.4	1.4
69	700	0.2	0.6
70	130	0.2	0.4

Example 22: Tumor Necrosis Factor Assays

5

Cell Culture.

The cells used in the assay are the human monocytic line U-937 (ATCC CRL-1593). The cells are grown in RPMI w/10% FCS and PSG supplement (R-10) and are not permitted to overgrow. The assay is carried out as follows:

10

SUBSTITUTE SHEET (RULE 26)

-134-

1. Count, then harvest cells by centrifugation. Resuspend the pellet in R-10 supplement to a concentration of 1.540×10^6 cells/mL.
- 5 2. Add test compound in 65 μ L R-10 to the appropriate wells of a 96-well flat bottom tissue culture plate. The initial dilution from a DMSO stock (100 mM compound) provides a 400 μ M solution, from which five additional three-fold serial
10 dilutions are made. Each dilution of 65 μ L (in triplicate) yields final compound test concentrations of 100 μ M, 33.3 μ M, 11.1 μ M, 3.7 μ M, 1.2 μ M and 0.4 μ M.
3. The counted, washed and resuspended
15 cells (200,000 cells/well) in 130 μ L are added to the wells.
4. Incubation is for 45 minutes to one hour at 37°C in 5% CO₂ in a water saturated container.
5. R-10 (65 μ L) containing 160 ng/mL PMA
20 (Sigma) is added to each well.
6. The test system is incubated at 37°C in 5% CO₂ overnight (18-20 hours) under 100% humidity.
7. Supernatant, 150 μ L, is carefully removed from each well for use in the ELISA assay.
- 25 8. For toxicity, a 50 μ L aliquot of working solution containing 5 mL R-10, 5 mL MTS solution [CellTiter 96 Aqueous One Solution Cell Proliferation Assay Cat.#G358/0,1 (Promega Biotech)] and 250 μ L PMS solution are added to each well containing the
30 remaining supernatant and cells and the cells incubated at 37°C in 5% CO₂ until the color develops. The system is excited at 570 nm and read at 630 nm.

SUBSTITUTE SHEET (RULE 26)

TNF α ELISA Assay

Coat Immulon[®] 2 plates with 0.1 mL/well of
1 μ g/mL Genzyme mAb in 0.1 M NaHCO₃ pH 8.0 buffer
5 overnight (about 18-20 hours) at 4°C, wrapped tightly
in Saran[®] wrap.

Flick out coating solution and block plates
with 0.3 mL/well blocking buffer overnight at 4°C,
wrapped in Saran[®] wrap.

10 Wash wells thoroughly 4X with wash buffer
and completely remove all wash buffer. Add 0.1
mL/well of either samples or rhTNF α standards.
Dilute samples if necessary in appropriate diluent
(e.g. tissue culture medium). Dilute standard in
15 same diluent. Standards and samples should be in
triplicates.

Incubate at 37°C for 1 hour in humidified
container.

Wash plates as above. Add 0.1 mL/well of
20 1:200 dilution of Genzyme rabbit anti-hTNF α .

Repeat incubation.

Repeat wash. Add 0.1 mL/well of 1 μ g/mL
Jackson goat anti-rabbit IgG (H+L)-peroxidase.

Incubate at 37°C for 30 minutes.

25 Repeat wash. Add 0.1 mL/well of peroxide-
ABTS solution.

Incubate at room temperature for 5-20
minutes.

Read OD at 405 nm.

30 12 Reagents are:

Genzyme mouse anti-human TNF monoclonal
(Cat.# 80-3399-01)

-136-

Genzyme rabbit anti-human TNF polyclonal
(Cat.#IP-300)

Genzyme recombinant human TNF (Cat.#TNF-H).

Jackson ImmunoResearch peroxide-conjugated
5 goat anti-rabbit IgG (H+L) (Cat.#111-035-144).

Kirkegaard/Perry peroxide ABTS solution
(Cat#50-66-01).

Immulon 2 96-well microtiter plates.

Blocking solution is 1 mg/mL gelatin in PBS
10 with 1X thimerasol.

Wash buffer is 0.5 mL Tween[®] 20 in 1 liter
of PBS.

15

20

Results:

25

Table 2

Inhibition of TNF? Release

30

Example Number	Per Cent Inhibition of TNF? Release in Micromolar (μ M) or IC ₅₀ (μ M)
----------------	--


SUBSTITUTE SHEET (RULE 26)

	[average]
3	3.72% @ 10 μ M (n=5)
7	IC ₅₀ = >50 μ M
4	-3.8% @10 (n=3) IC ₅₀ = >50 μ M
8	IC ₅₀ = >50 μ M
22	IC ₅₀ = >29 μ M
5	-10.8% @ 10 (n=3) IC ₅₀ = >50 μ M
23	IC ₅₀ = >50 μ M
24	IC ₅₀ = >50 μ M
25	IC ₅₀ = 14.0 μ M

From the foregoing, it will be observed
5 that numerous modifications and variations can be
effectuated without departing from the true spirit
and scope of the novel concepts of the present
invention. It is to be understood that no limitation
with respect to the specific example presented is
10 intended or should be inferred. The disclosure is
intended to cover by the appended claims all such
modifications as fall within the scope of the claims.

What is Claimed:

1. A compound corresponding to Formula A, or a pharmaceutically acceptable salt thereof:



5

A

wherein

R^2 is selected from the group consisting of a hydrido, C_1 - C_4 hydrocarbyl, hydroxy- C_1 - C_4 hydrocarbyl, C_1 - C_4 hydrocarbyloxy, halo- C_1 - C_4 hydrocarbyl, C_1 - C_4 hydrocarbyloxymethyl, aryl, aryl C_1 - C_4 hydrocarbyl, aminomethyl, (N- C_1 - C_3 hydrocarbyl)aminomethyl, (N,N-di- C_1 - C_3 hydrocarbyl)aminomethyl, (N-morpholino)methyl, (N-pyrrolidino)methyl and a (N-thiomorpholino)methyl group;

R^1 is a substituent that contains an optionally substituted 5- or 6-membered carbocyclo, heterocyclo, aryl or heteroaryl radical bonded directly to the depicted SO_2 -group that has a length greater than that of an extended pentyl group and less than about that of an extended eicosyl group, wherein R^1 defines a three-dimensional volume upon rotation about an axis drawn through the SO_2 -bonded 1-position and the 4-position of a 6-membered ring radical or drawn through the SO_2 -bonded 1-position and the center of 3,4-bond of a 5-membered ring

radical, the widest dimension of said volume in a direction transverse to the axis of rotation being about that of one furanyl ring to about that of two phenyl rings;

- 5 R⁶ and R⁷ are independently selected from the group consisting of a hydrido, alkylcarbonyl, arylcarbonyl, alkoxy carbonyl, alkylaminocarbonyl, aralkyl, carboxyalkyl, carboxyalkylcarbonyl, alkoxy carbonylaminoalkylcarbonyl,
- 10 heterocyclocarbonyl, heterocycloalkylcarbonyl, aryloxy carbonylaminoalkylcarbonyl, aralkyloxy carbonylaminoalkylcarbonyl, perfluoroalkylcarbonyl, trifluoromethylalkylcarbonyl, alkylthioalkylcarbonyl, arylthioalkylcarbonyl,
- 15 aralkylthioalkylcarbonyl, heteroaralkylthioalkylcarbonyl, or a sulfoxide or sulfone of any of said thio substituents, alkylsulfonyl, arylsulfonyl and an aminocarbonylalkyl or aminocarbonylalkylcarbonyl group
- 20 wherein the aminocarbonylalkyl or aminocarbonylalkylcarbonyl nitrogen is (i) unsubstituted, or (ii) is the reacted amine of an amino acid, or (iii) substituted with one or two radicals selected from the group consisting
- 25 of an alkyl, hydroxyalkyl, hydroxyheteroaralkyl, cycloalkyl, aralkyl, trifluoromethylalkyl, heterocycloalkyl, benzofused heterocycloalkyl, benzofused heterocycloalkyl, benzofused cycloalkyl, and an N,N-dialkylsubstituted-
- 30 alkylaminoalkyl group, or (iv) that carboxamido nitrogen and two substituents bonded thereto together form a 5- to 8-membered heterocyclo, heteroaryl or benzofused heterocycloalkyl ring that is itself unsubstituted or substituted with

one or two radicals independently selected from the group consisting of an alkyl, alkoxy carbonyl, nitro, heterocycloalkyl, hydroxy, hydroxycarbonyl, aryl, aralkyl, heteroaralkyl and an amino group, or

R^6 and R^7 together with the depicted nitrogen atom form a saturated or unsaturated cyclic imide substituent in which the ring contains five to eight atoms; and

R^{20} is (a) $-O-R^{21}$, where R^{21} is selected from the group consisting of a hydrido, C_1-C_6 -alkyl, aryl, ar- C_1-C_6 -alkyl group and a pharmaceutically acceptable cation, (b) $-NR^{13}-O-R^{22}$ wherein R^{22} is a selectively removable protecting group and R^{13} is a hydrido, C_1-C_6 -alkyl or benzyl group, or (c) $-NR^{13}-O-R^{14}$, wherein R^{14} is a hydrido, a pharmaceutically acceptable cation or $C(W)R^{15}$ group, where W is O or S and R^{15} is selected from the group consisting of an C_1-C_6 -alkyl, aryl, C_1-C_6 -alkoxy, heteroaryl- C_1-C_6 -alkyl, C_3-C_8 -cycloalkyl- C_1-C_6 -alkyl, aryloxy, ar- C_1-C_6 -alkoxy, ar- C_1-C_6 -alkyl, heteroaryl and amino C_1-C_6 -alkyl group wherein the aminoalkyl nitrogen is (i) unsubstituted or (ii) substituted with one or two substituents independently selected from the group consisting of an C_1-C_6 -alkyl, aryl, ar- C_1-C_6 -alkyl, C_3-C_8 -cycloalkyl- C_1-C_6 -alkyl, ar- C_1-C_6 -alkoxy carbonyl, C_1-C_6 -alkoxy carbonyl, and C_1-C_6 -alkanoyl radical, or (iii) wherein the amino C_1-C_6 -alkyl nitrogen and two substituents attached thereto form a 5- to 8-membered heterocyclo or heteroaryl ring.

2. The compound or salt according to claim 1 wherein R^1 has length that is longer than an extended hexyl group and less than that of an extended stearyl group.

3. The compound or salt according to claim 1 wherein the 5- or 6-membered carbocyclo, heterocyclo, aryl or heteroaryl radical of R^1 is substituted with an -A-R-E-Y substituent in which A is selected from the group consisting of

- (1) -O-;
- (2) -S-;
- (3) -NR^k-;
- (4) -CO-N(R^k) or -N(R^k)-CO-;
- (5) -CO-O- or -O-CO-;
- (6) -O-CO-O-;
- (7) -HC=CH-;
- (8) -NH-CO-NH-;
- (9) -C≡C-;
- (10) -N=N-;
- (11) -NH-NH-;
- (12) -CS-N(R^k) - or -N(R^k)-CS-;
- (13) -CH₂-;
- (14) -O-CH₂- or -CH₂-O-;
- (15) -S-CH₂- or -CH₂-S-;
- (16) -SO-; and
- (17) -SO₂-; or
- (18) A is absent and R is directly bonded to the depicted SO₂-bonded 5- or 6-membered ring group;

the moiety R is selected from the group consisting of alkyl, alkoxy, halo, alkoxyalkyl, aryl, heteroaryl, cycloalkyl, heterocycloalkyl, aralkyl, heteroaralkyl, heterocycloalkyl, cycloalkyl, 5 cycloalkoxyalkyl, heterocycloalkoxyalkyl, aryloxyalkyl, heteroaryloxy-alkyl, arylthioalkyl, heteroarylthioalkyl, cycloalkylthioalkyl, and a heterocycloalkylthioalkyl group wherein the aryl, heteroaryl, cycloalkyl or heterocycloalkyl 10 substituent is (i) unsubstituted or (ii) substituted with one or two radicals selected from the group consisting of a halo, alkyl, perfluoroalkyl, perfluoroalkoxy, perfluoroalkylthio, trifluoromethylalkyl, amino, alkoxycarbonylalkyl, 15 alkoxy, C₁-C₂-alkylenedioxy, hydroxycarbonylalkyl, hydroxycarbonylalkylamino, nitro, hydroxy, hydroxyalkyl, alkanoylamino, and a alkoxycarbonyl group;

the group E is selected from the group 20 consisting of

- (1) -CO(R^g)- or -(R^g)CO-;
- (2) -CON(R^k)- or -(R^k)NCO-;
- (3) -CO-;
- (4) -SO₂-R^g or -R^gSO₂-;
- 25 (5) -SO₂-;
- (6) -N(R^k)-SO₂- or -SO₂-N(R^k)-; or
- (7) E is absent and R is bonded directly to Y; and

Y is absent or is selected from the group 30 consisting of a hydrido, alkyl, alkoxy, haloalkyl, aryl, aralkyl, cycloalkyl, heteroaryl, hydroxy, nitrile, nitro, aryloxy, aralkoxy, heteroaryloxy, heteroaralkyl, R^coxyalkyl, perfluoroalkoxy,

perfluoroalkylthio, trifluoromethylalkyl, alkenyl,
heterocycloalkyl, cycloalkyl, trifluoromethyl,
alkoxycarbonyl, and a aminoalkyl group, wherein the
aryl, heteroaryl or heterocycloalkyl group is (i)
5 unsubstituted or (ii) substituted with one or two
radicals independently selected from the group
consisting of an alkanoyl, halo, nitro, nitrile,
haloalkyl, alkyl, aralkyl, aryl, alkoxy, and an amino
group wherein the amino nitrogen is (i) unsubstituted
10 or (ii) substituted with one or two groups
independently selected from hydrido, alkyl, and an
aralkyl group; and wherein

R^C is selected from the group consisting of
a hydrido, alkanoyl, arylalkyl, aroyl,
15 bisalkoxyalkyl, alkyl, haloalkyl, perfluoroalkyl,
trifluoromethylalkyl, perfluoroalkoxyalkyl,
alkoxyalkyl, cycloalkyl, heterocycloalkyl,
heterocycloalkylcarbonyl, aryl, heterocyclo,
heteroaryl, cycloalkyl, aryloxyalkyl,
20 heteroaryloxyalkyl, heteroarylalkoxyalkyl,
heteroarylthioalkyl, arylsulfonyl, alkylsulfonyl,
heteroarylsulfonyl, carboxyalkyl,
alkoxycarbonylalkyl, aminocarbonyl,
alkyliminocarbonyl, aryliminocarbonyl,
25 heterocycloiminocarbonyl, arylthioalkyl,
alkylthioalkyl, arylthioalkenyl, alkylthioalkenyl,
heteroarylalkyl, haloalkanoyl, hydroxyalkanoyl,
thiolalkanoyl, alkenyl, alkynyl, alkoxyalkyl,
alkoxycarbonyl, aryloxycarbonyl, R^{dRe} aminoalkyl-
30 carbonyl, hydroxyalkyl, R^{dRe} aminoalkyl,
 R^{dRe} aminoalkylsulfonyl, R^{dRe} aminosulfonyl wherein the
amino nitrogen is (i) unsubstituted or (ii)
independently substituted with one or two R^d
radicals, or the substituents on the amino group

taken together with the amino nitrogen form a saturated or partially unsaturated heterocyclo group optionally substituted with one, two or three groups independently selected from R^d substituents or a
5 heteroaryl group optionally substituted with one, two or three groups independently selected from R^f substituents;

R^d and R^e are independently selected from the group consisting of a hydrido, alkyl, alkenyl,
10 alkenyl, arylalkyl, aryl, alkanoyl, aroyl, arylalkylcarbonyl, alkoxycarbonyl or arylalkyloxycarbonyl group;

R^f is selected from the group consisting of a nitro, hydroxy, alkyl, halogen, aryl, alkoxy,
15 cyano, and R^dR^e amino group;

R^g is selected from the group consisting of a hydrido, aryl, heteroaryl, heterocyclo, aroyl, alkanoyl, heteroaroyl, halogen, cyano, aldehydo, hydroxy, amino, alkyl, alkenyl, alkynyl, cycloalkyl,
20 cycloalkenyl, alkoxy, aryloxy, heteroaryloxy, alkenyloxy, alkynyloxy, alkoxyaryl, alkoxyheteroaryl, R^{hR^i} -amino, alkoxyalkyl, alkylenedioxy, aryloxyalkyl, perfluoroalkyl, trifluoroalkyl, alkylthio, arylthio, alkyloxycarbonyl, alkyloxycarbonyloxy,
25 aryloxycarbonyl, arylalkyloxycarbonyl, arylalkyloxycarbonylamino, aryloxycarbonyloxy, carboxy, R^{hR^i} -aminocarbonyloxy, R^{hR^i} -aminocarbonyl, R^{hR^i} -aminoalkanoyl, hydroxyaminocarbonyl, R^{hR^i} -aminosulfonyl, R^{hR^i} -aminocarbonyl (R^h) amino,
30 trifluoromethylsulfonyl- (R^h) amino, heteroarylsulfonyl (R^h) amino, arylsulfonyl (R^h) amino, arylsulfonyl (R^h) aminocarbonyl,

alkylsulfonyl (R^h) amino,
arylcarbonyl (R^h) aminosulfonyl, and an
alkylsulfonyl (R^h) aminocarbonyl substituent;

wherein R^h is selected from the group
5 consisting of an arylalkyl, aryl, heteroaryl,
heterocyclo, alkyl, alkynyl, alkenyl, alkoxyalkyl,
substituted or unsubstituted aminoalkyl,
alkyloxycarbonyl, arylalkyloxycarbonyl, carboxyalkyl,
haloalkyl, alkanoyl, aroyl, substituted or
10 unsubstituted aminoalkanoyl, halo alkanoyl and a
hydroxyalkyl group, each of which groups is
optionally substituted by one or two groups
independently selected from R^j substituents as are
the substituents of the substituted aminoalkyl and
15 substituted aminoalkanoyl groups;

R^i is selected from the group consisting of
an arylalkyl, aryl, heteroaryl, heterocyclo, alkyl,
alkynyl, alkenyl, alkoxyalkyl, alkoxyalkyl,
substituted or unsubstituted aminoalkyl,
20 alkyloxycarbonyl, arylalkyloxycarbonyl, carboxyalkyl,
haloalkyl, alkanoyl, aroyl, substituted or
unsubstituted aminoalkanoyl, halo alkanoyl and a
hydroxyalkyl group, each of which groups are
optionally substituted by one or two R^j substituents;

25 R^j is selected from the group consisting of
an arylalkyl, aryl, heteroaryl, heterocyclo, alkyl,
alkynyl, alkenyl, alkoxyalkyl, alkoxyalkyl,
substituted or unsubstituted aminoalkyl,
alkyloxycarbonyl, arylalkyloxycarbonyl, carboxyalkyl,
30 haloalkyl, alkanoyl, aroyl, substituted or
unsubstituted aminoalkanoyl, halo alkanoyl and a
hydroxyalkyl group, wherein the substituents of the
substituted aminoalkyl and substituted aminoalkanoyl

groups are selected from the group consisting of an alkyl, alkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, aryloxy carbonyl and an alkyloxy carbonyl group; and

- 5 R^k is selected from hydrido, alkyl, alkenyl, alkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, aryloxy carbonyl, alkyloxy carbonyl, R^{CR^d} amino carbonyl, R^{CR^d} aminosulfonyl, R^{CR^d} aminoalkanoyl and R^{CR^d} aminoalkylsulfonyl.

10

4. The compound or salt according to claim 3 wherein R^1 is an aryl or heteroaryl group that is substituted by said -A-R-E-Y substituent.

15

5. The compound or salt according to claim 1 wherein R^2 is selected from the group consisting of a hydrido, hydroxy, hydroxymethyl, methoxymethyl, C_1 - C_4 hydrocarbyl, aryl and a methyl-N-morpholinyl group.


20

6. The compound or salt according to claim 5 wherein said R^1 substituent is PhR^4 in which R^4 is bonded at the 4-position of the SO_2 -linked phenyl (Ph) radical, and R^4 is selected from the group
- 25 consisting of a hydrocarbyl or hydrocarbyloxy group, a phenyl group, a phenoxy group, a thiophenoxy group, an anilino group, a phenylazo group, a phenylureido group, a benzamido group, a nicotinamido group, an isonicotinamido group, and a picolinamido group..


30

7. The compound or salt according to claim 6 wherein said compound corresponds in structure to the formula below

SUBSTITUTE SHEET (RULE 26)




8. The compound or salt according to claim 6 wherein said compound corresponds in structure to the formula below




5

9. The compound or salt according to claim 6 wherein said compound corresponds in structure to the formula below




10


10. The compound or salt according to claim 6 wherein said compound corresponds in structure to the formula below



11. The compound or salt according to claim 6 wherein said compound corresponds in structure to the formula below




12. The compound or salt according to claim 6 wherein said compound corresponds in structure to the formula below




5

13. The compound or salt according to claim 6 wherein said compound corresponds in structure to the formula below




10

14. The compound or salt according to claim 6 wherein said compound corresponds in structure to the formula below




15. The compound or salt according to claim 6 wherein said compound corresponds in structure to the formula below

-149-




16. The compound or salt according to claim 6 wherein said compound corresponds in structure to the formula below




5

17. The compound or salt according to claim 6 wherein said compound corresponds in structure to the formula below




10

18. The compound or salt according to claim 6 wherein said compound corresponds in structure to the formula below




15

19. The compound or salt according to claim 6 wherein said compound corresponds in structure to the formula below




SUBSTITUTE SHEET (RULE 26)


20. The compound or salt according to claim 6 wherein said compound corresponds in structure to the formula below




5 21. The compound or salt according to claim 6 wherein said compound corresponds in structure to the formula below




10 22. The compound or salt according to claim 6 wherein said compound corresponds in structure to the formula below




15 23. The compound or salt according to claim 6 wherein said compound corresponds in structure to the formula below




24. The compound or salt according to claim 6 wherein said compound corresponds in structure to the formula below




5 25. The compound or salt according to claim 6 wherein said compound corresponds in structure to the formula below




10 26. The compound or salt according to claim 6 wherein said compound corresponds in structure to the formula below




15 27. The compound or salt according to claim 6 wherein said compound corresponds in structure to the formula below



28. The compound or salt according to claim 6 wherein said compound corresponds in structure to the formula below




29. The compound or salt according to claim 6 wherein said compound corresponds in structure to the formula below



5

30. The compound or salt according to claim 6 wherein said compound corresponds in structure to the formula below



10


31. The compound or salt according to claim 1 wherein R^6 is hydrido, and R^7 is an aminoalkylcarbonyl or a substituted aminoalkylcarbonyl group.

15

32. The compound or salt according to claim 6 wherein R^7 is selected from the group consisting of an alkylcarbonyl, alkoxycarbonyl, arylcarbonyl, alkylsulfonyl, arylsulfonyl, alkoxycarbonylaminoalkylcarbonyl, cycloalkoxycarbonylaminoalkylcarbonyl, aryloxycarbonylaminoalkylcarbonyl, and heteroaryloxycarbonylaminoalkylcarbonyl group.

20

33. A compound corresponding to Formula B1, or a pharmaceutically acceptable salt thereof:



B1

5

wherein :

R^1 is a substituent that contains an optionally substituted 5- or 6-membered carbocyclo, heterocyclo, aryl or heteroaryl radical bonded
 10 directly to the depicted SO_2 -group that has a length greater than that of an extended pentyl group and less than about that of an extended eicosyl group, wherein R^1 defines a three-dimensional volume upon rotation about an axis drawn through the SO_2 -bonded
 15 1-position and the 4-position of a 6-membered ring radical or drawn through the SO_2 -bonded 1-position and the center of 3,4-bond of a 5-membered ring radical, the widest dimension of said volume in a direction transverse to the axis of rotation being
 20 about that of one furanyl ring to about that of two phenyl rings;

R^{7A} is selected from the group consisting of an alkylcarbonyl, alkoxycarbonyl, arylcarbonyl, alkylsulfonyl, arylsulfonyl, aminoalkylcarbonyl,
 25 aminoaralkylcarbonyl, alkoxycarbonylaminoalkylcarbonyl, alkoxycarbonylaminoaralkylcarbonyl, cycloalkoxycarbonylaminoalkylcarbonyl, cycloalkoxycarbonylaminoaralkylcarbonyl,

aryloxycarbonylaminoalkylcarbonyl, aryloxycarbonyl-
aminoaralkylcarbonyl, aralkyloxycarbonylamino-
alkylcarbonyl, aralkyloxycarbonylaminoalkylcarbonyl,
heteroaryloxycarbonylaminoalkylcarbonyl and a
5 heteroaryloxycarbonylaminoaralkylcarbonyl group; and
R¹⁴ is a hydrido, a pharmaceutically
acceptable cation or C(W)R¹⁵ group, where W is O or S
and R¹⁵ is selected from the group consisting of an
C₁-C₆-alkyl, aryl, C₁-C₆-alkoxy, heteroaryl-C₁-C₆-
10 alkyl, C₃-C₈-cycloalkyl-C₁-C₆-alkyl, aryloxy, ar-C₁-
C₆-alkoxy, ar-C₁-C₆-alkyl, heteroaryl and amino C₁-
C₆-alkyl group wherein the aminoalkyl nitrogen is (i)
unsubstituted or (ii) substituted with one or two
substituents independently selected from the group
15 consisting of an C₁-C₆-alkyl, aryl, ar-C₁-C₆-alkyl,
C₃-C₈-cycloalkyl-C₁-C₆-alkyl, ar-C₁-C₆-
alkoxycarbonyl, C₁-C₆-alkoxycarbonyl, and C₁-C₆-
alkanoyl radical, or (iii) wherein the amino C₁-C₆-
alkyl nitrogen and two substituents attached thereto
20 form a 5- to 8-membered heterocyclo or heteroaryl
ring.

34. The compound or salt according to
claim 33 wherein R¹ has length that is longer than an
25 extended hexyl group and less than that of an
extended stearyl group.

35. The compound or salt according to
claim 33 wherein the 5- or 6-membered carbocyclo,
30 heterocyclo, aryl or heteroaryl radical of R¹ is
substituted with an -A-R-E-Y substituent in which A
is selected from the group consisting of

- (1) -O-;
- (2) -S-;
- (3) -NR^k-;
- (4) -CO-N(R^k) or -N(R^k)-CO-;
- 5 (5) -CO-O- or -O-CO-;
- (6) -O-CO-O-;
- (7) -HC=CH-;
- (8) -NH-CO-NH-;
- (9) -C≡C-;
- 10 (10) -N=N-;
- (11) -NH-NH-;
- (12) -CS-N(R^k)- or -N(R^k)-CS-;
- (13) -CH₂-;
- (14) -O-CH₂- or -CH₂-O-;
- 15 (15) -S-CH₂- or -CH₂-S-;
- (16) -SO-; and
- (17) -SO₂-; or
- (18) A is absent and R is directly bonded to the depicted SO₂-bonded 5- or 6-membered
- 20 ring group;
- the moiety R is selected from the group consisting of alkyl, alkoxy, halo, alkoxyalkyl, aryl, heteroaryl, cycloalkyl, heterocycloalkyl, aralkyl, heteroaralkyl, heterocycloalkyl, cycloalkyl,
- 25 cycloalkoxyalkyl, heterocycloalkoxyalkyl, aryloxyalkyl, heteroaryloxy-alkyl, arylthioalkyl, heteroarylthioalkyl, cycloalkylthioalkyl, and a heterocycloalkylthioalkyl group wherein the aryl, heteroaryl, cycloalkyl or heterocycloalkyl
- 30 substituent is (i) unsubstituted or (ii) substituted with one or two radicals selected from the group consisting of a halo, alkyl, perfluoroalkyl, perfluoroalkoxy, perfluoroalkylthio,

trifluoromethylalkyl, amino, alkoxycarbonylalkyl,
alkoxy, C₁-C₂-alkylenedioxy, hydroxycarbonylalkyl,
hydroxycarbonylalkylamino, nitro, hydroxy,
hydroxyalkyl, alkanoylamino, and a alkoxycarbonyl
5 group;

the group E is selected from the group
consisting of

- (1) -CO(R^g) - or -(R^g)CO-;
- (2) -CON(R^k) - or -(R^k)NCO-;
- 10 (3) -CO-;
- (4) -SO₂-R^g or -R^gSO₂-;
- (5) -SO₂-;
- (6) -N(R^k)-SO₂- or -SO₂-N(R^k)-; or
- (7) E is absent and R is bonded directly
15 to Y; and

Y is absent or is selected from the group
consisting of a hydrido, alkyl, alkoxy, haloalkyl,
aryl, aralkyl, cycloalkyl, heteroaryl, hydroxy,
nitrile, nitro, aryloxy, aralkoxy, heteroaryloxy,
20 heteroaralkyl, R^coxyalkyl, perfluoroalkoxy,
perfluoroalkylthio, trifluoromethylalkyl, alkenyl,
heterocycloalkyl, cycloalkyl, trifluoromethyl,
alkoxycarbonyl, and a aminoalkyl group, wherein the
aryl, heteroaryl or heterocycloalkyl group is (i)
25 unsubstituted or (ii) substituted with one or two
radicals independently selected from the group
consisting of an alkanoyl, halo, nitro, nitrile,
haloalkyl, alkyl, aralkyl, aryl, alkoxy, and an amino
group wherein the amino nitrogen is (i) unsubstituted
30 or (ii) substituted with one or two groups
independently selected from hydrido, alkyl, and an
aralkyl group; and wherein

- R^C is selected from the group consisting of a hydrido, alkanoyl, arylalkyl, aroyl, bisalkoxyalkyl, alkyl, haloalkyl, perfluoroalkyl, trifluoromethylalkyl, perfluoroalkoxyalkyl, alkoxyalkyl, cycloalkyl, heterocycloalkyl, heterocycloalkylcarbonyl, aryl, heterocyclo, heteroaryl, cycloalkyl, aryloxyalkyl, heteroaryloxyalkyl, heteroarylalkoxyalkyl, heteroarylthioalkyl, arylsulfonyl, alkylsulfonyl, heteroarylsulfonyl, carboxyalkyl, alkoxycarbonylalkyl, aminocarbonyl, alkyliminocarbonyl, aryliminocarbonyl, heterocycloiminocarbonyl, arylthioalkyl, alkylthioalkyl, arylthioalkenyl, alkylthioalkenyl, heteroarylalkyl, haloalkanoyl, hydroxyalkanoyl, thiolalkanoyl, alkenyl, alkynyl, alkoxyalkyl, alkoxycarbonyl, aryloxycarbonyl, $R^d R^e$ aminoalkyl-carbonyl, hydroxyalkyl, $R^d R^e$ aminoalkyl, $R^d R^e$ aminoalkylsulfonyl, $R^d R^e$ aminosulfonyl wherein the amino nitrogen is (i) unsubstituted or (ii) independently substituted with one or two R^d radicals, or the substituents on the amino group taken together with the amino nitrogen form a saturated or partially unsaturated heterocyclo group optionally substituted with one, two or three groups independently selected from R^d substituents or a heteroaryl group optionally substituted with one, two or three groups independently selected from R^f substituents;
- R^d and R^e are independently selected from the group consisting of a hydrido, alkyl, alkenyl, alkenyl, arylalkyl, aryl, alkanoyl, aroyl,

arylalkylcarbonyl, alkoxycarbonyl or
arylalkyloxycarbonyl group;

R^f is selected from the group consisting of
a nitro, hydroxy, alkyl, halogen, aryl, alkoxy,
5 cyano, and R^{dRe} amino group;

R^g is selected from the group consisting of
a hydrido, aryl, heteroaryl, heterocyclo, aroyl,
alkanoyl, heteroaroyl, halogen, cyano, aldehydo,
hydroxy, amino, alkyl, alkenyl, alkynyl, cycloalkyl,
10 cycloalkenyl, alkoxy, aryloxy, heteroaryloxy,
alkenyloxy, alkynyloxy, alkoxyaryl, alkoxyheteroaryl,
 R^{hRi} -amino, alkoxyalkyl, alkylenedioxy, aryloxyalkyl,
perfluoroalkyl, trifluoroalkyl, alkylthio, arylthio,
alkyloxycarbonyl, alkyloxycarbonyloxy,
15 aryloxycarbonyl, arylalkyloxycarbonyl,
arylalkyloxycarbonylamino, aryloxycarbonyloxy,
carboxy, R^{hRi} -aminocarbonyloxy, R^{hRi} -aminocarbonyl,
 R^{hRi} -aminoalkanoyl, hydroxyaminocarbonyl, R^{hRi} -
aminosulfonyl, R^{hRi} -aminocarbonyl(R^h) amino,
20 trifluoromethylsulfonyl-(R^h) amino,
heteroarylsulfonyl(R^h) amino, arylsulfonyl(R^h) amino,
arylsulfonyl(R^h) aminocarbonyl,
alkylsulfonyl(R^h) amino,
arylcarbonyl(R^h) aminosulfonyl, and an
25 alkylsulfonyl(R^h) aminocarbonyl substituent;

wherein R^h is selected from the group
consisting of an arylalkyl, aryl, heteroaryl,
heterocyclo, alkyl, alkynyl, alkenyl, alkoxyalkyl,
substituted or unsubstituted aminoalkyl,
30 alkyloxycarbonyl, arylalkyloxycarbonyl, carboxyalkyl,
haloalkyl, alkanoyl, aroyl, substituted or
unsubstituted aminoalkanoyl, halo alkanoyl and a

hydroxyalkyl group, each of which groups is optionally substituted by one or two groups independently selected from R^j substituents as are the substituents of the substituted aminoalkyl and substituted aminoalkanoyl groups;

R^i is selected from the group consisting of an arylalkyl, aryl, heteroaryl, heterocyclo, alkyl, alkynyl, alkenyl, alkoxyalkyl, alkoxyalkyl, substituted or unsubstituted aminoalkyl, alkyloxycarbonyl, arylalkyloxycarbonyl, carboxyalkyl, haloalkyl, alkanoyl, aroyl, substituted or unsubstituted aminoalkanoyl, halo alkanoyl and a hydroxyalkyl group, each of which groups are optionally substituted by one or two R^j substituents;

R^j is selected from the group consisting of an arylalkyl, aryl, heteroaryl, heterocyclo, alkyl, alkynyl, alkenyl, alkoxyalkyl, alkoxyalkyl, substituted or unsubstituted aminoalkyl, alkyloxycarbonyl, arylalkyloxycarbonyl, carboxyalkyl, haloalkyl, alkanoyl, aroyl, substituted or unsubstituted aminoalkanoyl, halo alkanoyl and a hydroxyalkyl group, wherein the substituents of the substituted aminoalkyl and substituted aminoalkanoyl groups are selected from the group consisting of an alkyl, alkenyl, alkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, aryloxycarbonyl and an alkyloxycarbonyl group; and

R^k is selected from hydrido, alkyl, alkenyl, alkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, aryloxycarbonyl, alkyloxycarbonyl, $R^C R^d$ amino carbonyl, $R^C R^d$ aminosulfonyl, $R^C R^d$ aminoalkanoyl and $R^C R^d$ aminoalkylsulfonyl.

36. The compound or salt according to claim 35 wherein R^1 is an aryl or heteroaryl group, G, that is substituted by said -A-R-E-Y substituent.

5 37. The compound or salt according to claim 33 wherein said R^1 substituent is selected from the group consisting of a 4-(phenyl)phenyl, 4-(4'-methoxyphenyl)-phenyl, 4-(phenoxy)phenyl, 4-(thiophenyl)phenyl, 4-(phenylazo)phenyl 4-
10 (phenylureido)phenyl, 4-(anilino)phenyl, 4-(nicotinamido)phenyl, 4-(isonicotinamido)phenyl, 4-(picolinamido)phenyl and 4-(benzamido)phenyl group.

38. The compound or salt according to
15 claim 33 wherein said R^1 substituent is PhR^4 in which R^4 is bonded at the 4-position of the SO_2 -linked phenyl (Ph) radical, and R^4 is a single-ringed cyclohydrocarbyl, heterocyclo, aryl or heteroaryl group or another substituent having a chain length of
20 3 to about 14 carbon atoms.

39. The compound or salt according to claim 38 wherein said R^4 substituent is selected from the group consisting of a hydrocarbyl or
25 hydrocarbyloxy group, a phenyl group, a phenoxy group, a thiophenoxy group, an anilino group, a phenylazo group, a phenylureido group, a benzamido group, a nicotinamido group, an isonicotinamido group, and a picolinamido group.

30

40. The compound or salt according to claim 39 wherein said R^4 substituent is itself substituted with one or more substituent radicals at

the meta- or para-position or both of a six-membered ring with a single atom or a substituent containing a longest chain of up to ten atoms, excluding hydrogen.

- 5 41. The compound or salt according to claim 40 wherein said R⁴ substituent is substituted with a radical selected from the group consisting of a halo, hydrocarbyl, hydrocarbyloxy, nitro, cyano, perfluorohydrocarbyl, trifluoromethyl-hydrocarbyl,
- 10 hydroxy, mercapto, hydroxycarbonyl, aryloxy, arylthio, arylamino, arylhydrocarbyl, aryl, heteroaryloxy, heteroarylthio, heteroarylamino, heteroarhydrocarbyl, hydrocarbyloxycarbonyl-hydrocarbyl, heterocyclooxy, hydroxycarbonyl-
- 15 hydrocarbyl, heterocyclothio, heterocycloamino, cyclohydrocarbyloxy, cyclohydrocarbylthio, cyclohydrocarbylamino, heteroarylhydrocarbyloxy, heteroarylhydrocarbylthio, heteroaryl-
- hydrocarbylamino, arylhydrocarbyloxy,
- 20 arylhydrocarbylthio, arylhydrocarbylamino, heterocyclic, heteroaryl, hydroxycarbonyl-hydrocarbyloxy, alkoxycarbonylalkoxy, hydrocarbyloyl, arylcarbonyl, arylhydrocarbyloyl, hydrocarbyloxy, arylhydrocarbyloxy, hydroxyhydrocarbyl, hydroxy-
- 25 hydrocarbyloxy, hydrocarbylthio, hydrocarbyloxy-hydrocarbylthio, hydrocarbyloxycarbonyl, hydroxycarbonylhydrocarbyloxy, hydrocarbyloxy-carbonylhydrocarbyl, hydrocarbylhydroxycarbonyl-
- hydrocarbylthio, hydrocarbyloxycarbonyl-
- 30 hydrocarbyloxy, hydrocarbyloxycarbonyl-hydrocarbylthio, amino, hydrocarbylcarbonylamino, arylcarbonylamino, cyclohydrocarbylcarbonylamino, heterocyclohydrocarbylcarbonylamino, arylhydrocarbylcarbonylamino, heteroaryl-

carbonylamino, heteroarylhydrocarbylcarbonylamino,
heterocyclohydrocarbyloxy, hydrocarbylsulfonylamino,
arylsulfonylamino, arylhydrocarbylsulfonylamino,
heteroarylsulfonylamino, heteroarylhydrocarbyl-
5 sulfonylamino, cyclohydrocarbylsulfonylamino,
heterocyclohydrocarbylsulfonylamino and N-
monosubstituted or N,N-disubstituted aminohydrocarbyl
group wherein the substituent(s) on the nitrogen are
selected from the group consisting of hydrocarbyl,
10 aryl, arylhydrocarbyl, cyclohydrocarbyl,
arylhydrocarbyloxycarbonyl, hydrocarbyloxycarbonyl,
and hydrocarboyl, or wherein the nitrogen and two
substituents attached thereto form a 5- to 8-membered
heterocyclic or heteroaryl ring group.

15

42. The compound or salt according to
claim 38 wherein said R⁴ substituent is selected from
the group consisting of a heterocyclo,
heterocyclohydrocarbyl, arylhydrocarbyl,
20 arylheterocyclohydrocarbyl, heteroarylhydrocarbyl,
heteroarylheterocyclohydrocarbyl,
arylhydrocarbyloxyhydrocarbyl, aryloxyhydrocarbyl,
hydrocarboylhydrocarbyl, arylhydrocarboylhydrocarbyl,
arylcarbonylhydrocarbyl, arylazoaryl,
25 arylhydrazinoaryl, hydrocarbylthiohydrocarbyl,
hydrocarbylthioaryl, arylthiohydrocarbyl,
heteroarylthiohydrocarbyl, hydrocarbylthio-
arylhydrocarbyl, arylhydrocarbylthiohydrocarbyl,
arylhydrocarbylthioaryl, arylhydrocarbylamino,
30 heteroarylhydrocarbylamino, and a heteroarylthio
group.

43. The compound or salt according to
claim 42 wherein said R⁴ substituent is itself

substituted with one or more substituent radicals at the meta- or para-position or both of a six-membered ring with a single atom or a substituent containing a longest chain of up to ten atoms, excluding hydrogen.

5


44. The compound or salt according to claim 43 wherein said R⁴ substituent is substituted with a radical selected from the group consisting of a halo, hydrocarbyl, hydrocarbyloxy, nitro, cyano, perfluorohydrocarbyl, trifluoromethyl-hydrocarbyl, hydroxy, mercapto, hydroxycarbonyl, aryloxy, arylthio, arylamino, arylhydrocarbyl, aryl, heteroaryloxy, heteroarylthio, heteroarylamino, heteroarhydrocarbyl, hydrocarbyloxycarbonyl-hydrocarbyl, heterocyclooxy, hydroxycarbonyl-hydrocarbyl, heterocyclothio, heterocycloamino, cyclohydrocarbyloxy, cyclohydrocarbylthio, cyclohydrocarbylamino, heteroarylhydrocarbyloxy, heteroarylhydrocarbylthio, heteroaryl-hydrocarbylamino, arylhydrocarbyloxy, arylhydrocarbylthio, arylhydrocarbylamino, heterocyclic, heteroaryl, hydroxycarbonyl-hydrocarbyloxy, alkoxycarbonylalkoxy, hydrocarbyloyl, arylcarbonyl, arylhydrocarbyloyl, hydrocarbyloxy, arylhydrocarbyloxy, hydroxyhydrocarbyl, hydroxyhydrocarbyloxy, hydrocarbylthio, hydrocarbyloxyhydrocarbylthio, hydrocarbyloxycarbonyl, hydroxycarbonylhydrocarbyloxy, hydrocarbyloxy-carbonylhydrocarbyl, hydrocarbylhydroxycarbonyl-hydrocarbylthio, hydrocarbyloxycarbonyl-hydrocarbyloxy, hydrocarbyloxycarbonyl-hydrocarbylthio, amino, hydrocarbylcarbonylamino, arylcarbonylamino, cyclohydrocarbylcarbonylamino, heterocyclohydrocarbylcarbonylamino,

arylhydrocarbylcarbonylamino, heteroaryl-
carbonylamino, heteroarylhydrocarbylcarbonylamino,
heterocyclohydrocarbyloxy, hydrocarbylsulfonylamino,
arylsulfonylamino, arylhydrocarbylsulfonylamino,
5 heteroarylsulfonylamino, heteroarylhydrocarbyl-
sulfonylamino, cyclohydrocarbylsulfonylamino,
heterocyclohydrocarbylsulfonylamino and N-
monosubstituted or N,N-disubstituted aminohydrocarbyl
group wherein the substituent(s) on the nitrogen are
10 selected from the group consisting of hydrocarbyl,
aryl, arylhydrocarbyl, cyclohydrocarbyl,
arylhydrocarbyloxycarbonyl, hydrocarbyloxycarbonyl,
and hydrocarboyl, or wherein the nitrogen and two
substituents attached thereto form a 5- to 8-membered
15 heterocyclic or heteroaryl ring group.

45. The compound or salt according to
claim 37 wherein said R¹ substituent is a phenoxy
phenyl group.


20

46. The compound or salt according to
claim 45 wherein said compound corresponds in
structure to the formula below




25 47. The compound or salt according to
claim 45 wherein said compound corresponds in
structure to the formula below

-165-




48. The compound or salt according to claim 45 wherein said compound corresponds in structure to the formula below




5

49. The compound or salt according to claim 45 wherein said compound corresponds in structure to the formula below




10


50. The compound or salt according to claim 45 wherein said compound corresponds in structure to the formula below




51. The compound or salt according to claim 45 wherein said compound corresponds in structure to the formula below



52. The compound or salt according to claim 45 wherein said compound corresponds in structure to the formula below




53. The compound or salt according to claim 45 wherein said compound corresponds in structure to the formula below




54. The compound or salt according to claim 45 wherein said compound corresponds in structure to the formula below

-167-




55. The compound or salt according to claim 45 wherein said compound corresponds in structure to the formula below



5


56. The compound or salt according to claim 45 wherein said compound corresponds in structure to the formula below




10

57. The compound or salt according to claim 45 wherein said compound corresponds in structure to the formula below

-168-




58. The compound or salt according to
claim 45 wherein said compound corresponds in
5 structure to the formula below



59. The compound or salt according to
claim 37 wherein said R¹ substituent is a
10 4-(thiophenyl)phenyl group.


60. The compound or salt according to
claim 59 wherein said compound corresponds in
structure to the formula below




15

61. The compound or salt according to
claim 59 wherein said compound corresponds in
structure to the formula below


-169-



62. The compound or salt according to
claim 59 wherein said compound corresponds in
5 structure to the formula below




63. The compound or salt according to
claim 59 wherein said compound corresponds in
structure to the formula below



10


64. The compound or salt according to
claim 59 wherein said compound corresponds in
structure to the formula below




15

65. The compound or salt according to
claim 59 wherein said compound corresponds in
structure to the formula below

SUBSTITUTE SHEET (RULE 26)




66. The compound or salt according to claim 59 wherein said compound corresponds in structure to the formula below




5

67. The compound or salt according to claim 59 wherein said compound corresponds in structure to the formula below




10


68. The compound or salt according to claim 59 wherein said compound corresponds in structure to the formula below




69. The compound or salt according to claim 59 wherein said compound corresponds in structure to the formula below




5 70. The compound or salt according to claim 59 wherein said compound corresponds in structure to the formula below




10 71. The compound or salt according to claim 59 wherein said compound corresponds in structure to the formula below




15 72. The compound or salt according to claim 59 wherein said compound corresponds in structure to the formula below



73. The compound or salt according to
claim 59 wherein said compound corresponds in
5 structure to the formula below




74. The compound or salt according to
claim 59 wherein said compound corresponds in
structure to the formula below




10

75. The compound or salt according to
claim 59 wherein said compound corresponds in
structure to the formula below

-173-




76. The compound or salt according to claim 59 wherein said compound corresponds in structure to the formula below




5

77. The compound or salt according to claim 37 wherein said R¹ substituent is a 4-(benzamido)phenyl group.

10 78. The compound or salt according to claim 77 wherein said compound corresponds in structure to the formula below




15 79. The compound or salt according to claim 77 wherein said compound corresponds in structure to the formula below




80. The compound or salt according to claim 35 wherein said R^1 substituent is an aryl group substituted with an -A-R-E-Y substituent in which A,
 5 E and Y are absent..

81. The compound or salt according to claim 80 wherein said compound corresponds in structure to the formula below




10

82. The compound or salt according to claim 80 wherein said compound corresponds in structure to the formula below




15


83. The compound or salt according to claim 80 wherein said compound corresponds in structure to the formula below




84. The compound or salt according to claim 80 wherein said compound corresponds in structure to the formula below




5 85. The compound or salt according to claim 80 wherein said compound corresponds in structure to the formula below



10 86. The compound or salt according to claim 80 wherein said compound corresponds in structure to the formula below




15 87. The compound or salt according to claim 80 wherein said compound corresponds in structure to the formula below




88. The compound or salt according to claim 80 wherein said compound corresponds in structure to the formula below

-176-




89. The compound or salt according to claim 80 wherein said compound corresponds in structure to the formula below




5

90. The compound or salt according to claim 80 wherein said compound corresponds in structure to the formula below




10

91. The compound or salt according to claim 80 wherein said compound corresponds in structure to the formula below




15

92. The compound or salt according to claim 80 wherein said compound corresponds in structure to the formula below




93. The compound or salt according to claim 80 wherein said compound corresponds in structure to the formula below



5

94. A process for treating a host mammal having a condition associated with pathological matrix metalloprotease (MMP) activity that comprises administering to a mammalian host having such a condition an MMP enzyme-inhibiting effective amount of a compound corresponding in structure to Formula B that selectively inhibits one or more MMPs, while exhibiting less activity against at least MMP-1, or a salt of such a compound



B

wherein

R^2 is selected from the group consisting of a hydrido, C_1 - C_4 hydrocarbyl, hydroxy- C_1 - C_4 hydrocarbyl, C_1 - C_4 hydrocarbyloxy, halo- C_1 - C_4 hydrocarbyl, C_1 - C_4 hydrocarbyloxymethyl, aryl, aryl
5 C_1 - C_4 hydrocarbyl, aminomethyl, (N- C_1 - C_3 hydrocarbyl)aminomethyl, (N,N-di- C_1 - C_3 hydrocarbyl)aminomethyl, (N-morpholino)methyl, (N-pyrrolidino)methyl, and a (N-thiomorpholino)methyl group;

10 R^1 is a substituent that contains an optionally substituted 5- or 6-membered carbocyclo, heterocyclo, aryl or heteroaryl radical bonded directly to the depicted SO_2 -group that has a length greater than that of an extended pentyl group and
15 less than about that of an extended eicosyl group, wherein R^1 defines a three-dimensional volume upon rotation about an axis drawn through the SO_2 -bonded 1-position and the 4-position of a 6-membered ring radical or drawn through the SO_2 -bonded 1-position
20 and the center of 3,4-bond of a 5-membered ring radical, the widest dimension of said volume in a direction transverse to the axis of rotation being about that of one furanyl ring to about that of two phenyl rings;

25 R^6 and R^7 are independently selected from the group consisting of a hydrido, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, alkylaminocarbonyl, aralkyl, carboxyalkyl, carboxyalkylcarbonyl, alkoxycarbonylaminoalkylcarbonyl,
30 heterocyclocarbonyl, heterocycloalkylcarbonyl, aryloxycarbonylaminoalkylcarbonyl, aralkyloxycarbonylaminoalkylcarbonyl,

perfluoroalkylcarbonyl, trifluoromethylalkylcarbonyl,
alkylthioalkylcarbonyl, arylthioalkylcarbonyl,
aralkylthioalkylcarbonyl, heteroaralkylthio-
alkylcarbonyl, or a sulfoxide or sulfone of any of
5 said thio substituents, alkylsulfonyl, arylsulfonyl
and an aminocarbonylalkyl or aminocarbonyl-
alkylcarbonyl group

wherein the aminocarbonylalkyl or
aminocarbonylalkylcarbonyl nitrogen is (i)
10 unsubstituted, or (ii) is the reacted amine of
an amino acid, or (iii) substituted with one or
two radicals selected from the group consisting
of an alkyl, hydroxyalkyl, hydroxyheteroaralkyl,
cycloalkyl, aralkyl, trifluoromethylalkyl,
15 heterocycloalkyl, benzofused heterocycloalkyl,
benzofused heterocycloalkyl, benzofused
cycloalkyl, and an N,N-dialkylsubstituted-
alkylaminoalkyl group, or (iv) that carboxamido
nitrogen and two substituents bonded thereto
20 together form a 5- to 8-membered heterocyclo,
heteroaryl or benzofused heterocycloalkyl ring
that is itself unsubstituted or substituted with
one or two radicals independently selected from
the group consisting of an alkyl,
25 alkoxycarbonyl, nitro, heterocycloalkyl,
hydroxy, hydroxycarbonyl, aryl, aralkyl,
heteroaralkyl and an amino group, or

R⁶ and R⁷ together with the depicted
nitrogen atom form a saturated or unsaturated cyclic
30 imide substituent in which the ring contains five to
eight atoms; and

R¹³ is a hydrido, C₁-C₆-alkyl or benzyl
group; and

R^{14} is a hydrido, a pharmaceutically acceptable cation or $C(W)R^{15}$ group, where W is O or S and R^{15} is selected from the group consisting of an C_1 - C_6 -alkyl, aryl, C_1 - C_6 -alkoxy, heteroaryl- C_1 - C_6 -alkyl, C_3 - C_8 -cycloalkyl- C_1 - C_6 -alkyl, aryloxy, ar- C_1 - C_6 -alkoxy, ar- C_1 - C_6 -alkyl, heteroaryl and amino C_1 - C_6 -alkyl group wherein the aminoalkyl nitrogen is (i) unsubstituted or (ii) substituted with one or two substituents independently selected from the group consisting of an C_1 - C_6 -alkyl, aryl, ar- C_1 - C_6 -alkyl, C_3 - C_8 -cycloalkyl- C_1 - C_6 -alkyl, ar- C_1 - C_6 -alkoxycarbonyl, C_1 - C_6 -alkoxycarbonyl, and C_1 - C_6 -alkanoyl radical, or (iii) wherein the amino C_1 - C_6 -alkyl nitrogen and two substituents attached thereto form a 5- to 8-membered heterocyclo or heteroaryl ring.

95. The process according to claim 84 wherein R^2 , R^6 , and R^{13} are hydrido.

20

96. The process according to claim 84 wherein R^7 is R^{7A} , and R^{7A} is selected from the group consisting of an alkylcarbonyl, alkoxycarbonyl, arylcarbonyl, alkylsulfonyl, arylsulfonyl, aminoalkylcarbonyl, aminoaralkylcarbonyl, alkoxycarbonylamino-alkylcarbonyl, alkoxycarbonylaminoaralkylcarbonyl, cycloalkoxycarbonylaminoalkylcarbonyl, cycloalkoxycarbonylaminoaralkylcarbonyl, aryloxycarbonylaminoalkylcarbonyl, aryloxycarbonylaminoaralkylcarbonyl, aralkyloxycarbonylamino-alkylcarbonyl, aralkyloxycarbonylaminoalkylcarbonyl,

30

heteroaryloxycarbonylaminoalkylcarbonyl and a
heteroaryloxycarbonylaminoaralkylcarbonyl group.

97. The process according to claim 84
5 wherein the 5- or 6-membered carbocyclo, heterocyclo,
aryl or heteroaryl radical of R^1 is substituted with
an -A-R-E-Y substituent in which A is selected from
the group consisting of

- (1) -O-;
 - 10 (2) -S-;
 - (3) -NR^k-;
 - (4) -CO-N(R^k) or -N(R^k)-CO-;
 - (5) -CO-O- or -O-CO-;
 - (6) -O-CO-O-;
 - 15 (7) -HC=CH-;
 - (8) -NH-CO-NH-;
 - (9) -C≡C-;
 - (10) -N=N-;
 - (11) -NH-NH-;
 - 20 (12) -CS-N(R^k) - or -N(R^k)-CS-;
 - (13) -CH₂-;
 - (14) -O-CH₂- or -CH₂-O-;
 - (15) -S-CH₂- or -CH₂-S-;
 - (16) -SO-; and
 - 25 (17) -SO₂-; or
 - (18) A is absent and R is directly bonded
to the depicted SO₂-bonded 5- or 6-membered
ring group;
- the moiety R is selected from the group
30 consisting of alkyl, alkoxy, halo, alkoxyalkyl, aryl,
heteroaryl, cycloalkyl, heterocycloalkyl, aralkyl,
heteroaralkyl, heterocycloalkyl, cycloalkyl,
cycloalkoxyalkyl, heterocycloalkoxyalkyl,

aryloxyalkyl, heteroaryloxy-alkyl, arylthioalkyl, heteroarylthioalkyl, cycloalkylthioalkyl, and a heterocycloalkylthioalkyl group wherein the aryl, heteroaryl, cycloalkyl or heterocycloalkyl

- 5 substituent is (i) unsubstituted or (ii) substituted with one or two radicals selected from the group consisting of a halo, alkyl, perfluoroalkyl, perfluoroalkoxy, perfluoroalkylthio, trifluoromethylalkyl, amino, alkoxycarbonylalkyl, 10 alkoxy, C₁-C₂-alkylenedioxy, hydroxycarbonylalkyl, hydroxycarbonylalkylamino, nitro, hydroxy, hydroxyalkyl, alkanoylamino, and a alkoxycarbonyl group;

the group E is selected from the group
15 consisting of

- (1) -CO(R^g) - or -(R^g)CO-;
- (2) -CON(R^k) - or -(R^k)NCO-;
- (3) -CO-;
- (4) -SO₂-R^g or -R^gSO₂-;
- 20 (5) -SO₂-;
- (6) -N(R^k)-SO₂- or -SO₂-N(R^k)-; or
- (7) E is absent and R is bonded directly to Y; and

the moiety Y is absent or is selected from
25 the group consisting of a hydrido, alkyl, alkoxy, haloalkyl, aryl, aralkyl, cycloalkyl, heteroaryl, hydroxy, nitrile, nitro, aryloxy, aralkoxy, heteroaryloxy, heteroaralkyl, R^coxyalkyl, perfluoroalkoxy, perfluoroalkylthio, 30 trifluoromethylalkyl, alkenyl, heterocycloalkyl, cycloalkyl, trifluoromethyl, alkoxycarbonyl, and a aminoalkyl group, wherein the aryl, heteroaryl or heterocycloalkyl group is (i) unsubstituted or (ii)

substituted with one or two radicals independently
selected from the group consisting of an alkanoyl,
halo, nitro, nitrile, haloalkyl, alkyl, aralkyl,
aryl, alkoxy, and an amino group wherein the amino
5 nitrogen is (i) unsubstituted or (ii) substituted
with one or two groups independently selected from
hydrido, alkyl, and an aralkyl group; and wherein

R^C is selected from the group consisting of
a hydrido, alkanoyl, arylalkyl, aroyl,
10 bisalkoxyalkyl, alkyl, haloalkyl, perfluoroalkyl,
trifluoromethylalkyl, perfluoroalkoxyalkyl,
alkoxyalkyl, cycloalkyl, heterocycloalkyl,
heterocycloalkylcarbonyl, aryl, heterocyclo,
heteroaryl, cycloalkyl, aryloxyalkyl,
15 heteroaryloxyalkyl, heteroarylalkoxyalkyl,
heteroarylthioalkyl, arylsulfonyl, alkylsulfonyl,
heteroarylsulfonyl, carboxyalkyl,
alkoxycarbonylalkyl, aminocarbonyl,
alkyliminocarbonyl, aryliminocarbonyl,
20 heterocycloiminocarbonyl, arylthioalkyl,
alkylthioalkyl, arylthioalkenyl, alkylthioalkenyl,
heteroarylalkyl, haloalkanoyl, hydroxyalkanoyl,
thiolalkanoyl, alkenyl, alkynyl, alkoxyalkyl,
alkoxycarbonyl, aryloxycarbonyl, R^{dRe} aminoalkyl-
25 carbonyl, hydroxyalkyl, R^{dRe} aminoalkyl,
 R^{dRe} aminoalkylsulfonyl, R^{dRe} aminosulfonyl wherein the
amino nitrogen is (i) unsubstituted or (ii)
independently substituted with one or two R^d
radicals, or the substituents on the amino group
30 taken together with the amino nitrogen form a
saturated or partially unsaturated heterocyclo group
optionally substituted with one, two or three groups
independently selected from R^d substituents or a

heteroaryl group optionally substituted with one, two or three groups independently selected from R^f substituents;

R^d and R^e are independently selected from
5 the group consisting of a hydrido, alkyl, alkenyl, alkenyl, arylalkyl, aryl, alkanoyl, aroyl, arylalkylcarbonyl, alkoxycarbonyl or arylalkylloxycarbonyl group;

R^f is selected from the group consisting of
10 a nitro, hydroxy, alkyl, halogen, aryl, alkoxy, cyano, and R^{dRe} amino group;

R^g is selected from the group consisting of
a hydrido, aryl, heteroaryl, heterocyclo, aroyl, alkanoyl, heteroaroyl, halogen, cyano, aldehydo,
15 hydroxy, amino, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, alkoxy, aryloxy, heteroaryloxy, alkenyloxy, alkynyloxy, alkoxyaryl, alkoxyheteroaryl, R^{hRi} -amino, alkoxyalkyl, alkylenedioxy, aryloxyalkyl, perfluoroalkyl, trifluoroalkyl, alkylthio, arylthio,
20 alkyloxycarbonyl, alkyloxycarbonyloxy, aryloxycarbonyl, arylalkylloxycarbonyl, arylalkylloxycarbonylamino, aryloxycarbonyloxy, carboxy, R^{hRi} -aminocarbonyloxy, R^{hRi} -aminocarbonyl, R^{hRi} -aminoalkanoyl, hydroxyaminocarbonyl, R^{hRi} -
25 aminosulfonyl, R^{hRi} -aminocarbonyl (R^h) amino, trifluoromethylsulfonyl- (R^h) amino, heteroarylsulfonyl (R^h) amino, arylsulfonyl (R^h) amino, arylsulfonyl (R^h) aminocarbonyl, alkylsulfonyl (R^h) amino,
30 arylcarbonyl (R^h) aminosulfonyl, and an alkylsulfonyl (R^h) aminocarbonyl substituent;

wherein R^h is selected from the group consisting of an arylalkyl, aryl, heteroaryl, heterocyclo, alkyl, alkynyl, alkenyl, alkoxyalkyl, substituted or unsubstituted aminoalkyl, alkyloxycarbonyl, arylalkyloxycarbonyl, carboxyalkyl, haloalkyl, alkanoyl, aroyl, substituted or unsubstituted aminoalkanoyl, halo alkanoyl and a hydroxyalkyl group, each of which groups is optionally substituted by one or two groups independently selected from R^j substituents as are the substituents of the substituted aminoalkyl and substituted aminoalkanoyl groups;

R^i is selected from the group consisting of an arylalkyl, aryl, heteroaryl, heterocyclo, alkyl, alkynyl, alkenyl, alkoxyalkyl, alkoxyalkyl, substituted or unsubstituted aminoalkyl, alkyloxycarbonyl, arylalkyloxycarbonyl, carboxyalkyl, haloalkyl, alkanoyl, aroyl, substituted or unsubstituted aminoalkanoyl, halo alkanoyl and a hydroxyalkyl group, each of which groups are optionally substituted by one or two R^j substituents;


R^j is selected from the group consisting of an arylalkyl, aryl, heteroaryl, heterocyclo, alkyl, alkynyl, alkenyl, alkoxyalkyl, alkoxyalkyl, substituted or unsubstituted aminoalkyl, alkyloxycarbonyl, arylalkyloxycarbonyl, carboxyalkyl, haloalkyl, alkanoyl, aroyl, substituted or unsubstituted aminoalkanoyl, halo alkanoyl and a hydroxyalkyl group, wherein the substituents of the substituted aminoalkyl and substituted aminoalkanoyl groups are selected from the group consisting of an alkyl, alkenyl, alkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, aryloxycarbonyl and an alkyloxycarbonyl group; and

R^k is selected from hydrido, alkyl, alkenyl, alkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, aryloxycarbonyl, alkyloxycarbonyl, $R^C R^d$ amino carbonyl, $R^C R^d$ aminosulfonyl, $R^C R^d$ aminoalkanoyl and $R^C R^d$ aminoalkylsulfonyl.

98. The process according to claim 84 wherein R^1 has length that is longer than an extended hexyl group and less than that of an extended stearyl group.

99. The process according to claim 84 wherein said compound or salt also does not substantially inhibit the production of TNF.

100. A compound corresponding in structure to formula C, below.



C

20

wherein

D is a nucleophilically-displacable leaving group;

R^2 is selected from the group consisting of a hydrido, C_1 - C_4 hydrocarbyl, hydroxy- C_1 - C_4 hydrocarbyl, C_1 - C_4 hydrocarbyloxy, halo- C_1 - C_4 hydrocarbyl, C_1 - C_4 hydrocarbyloxymethyl, aryl, aryl

C₁-C₄ hydrocarbyl, aminomethyl, (N-C₁-C₃
hydrocarbyl)aminomethyl, (N,N-di-C₁-C₃
hydrocarbyl)aminomethyl, (N-morpholino)methyl, (N-
pyrrolidino)methyl, and a (N-thiomorpholino)methyl
5 group;

R⁶ and R⁷ are independently selected from
the group consisting of a hydrido, alkylcarbonyl,
arylcarbonyl, alkoxycarbonyl, alkylaminocarbonyl,
aralkyl, carboxyalkyl, carboxyalkylcarbonyl,
10 alkoxycarbonylaminoalkylcarbonyl,
heterocyclocarbonyl, heterocycloalkylcarbonyl,
aryloxycarbonylaminoalkylcarbonyl,
aralkyloxycarbonylaminoalkylcarbonyl,
perfluoroalkylcarbonyl, trifluoromethylalkylcarbonyl,
15 alkylthioalkylcarbonyl, arylthioalkylcarbonyl,
aralkylthioalkylcarbonyl, heteroaralkylthio-
alkylcarbonyl, or a sulfoxide or sulfone of any of
said thio substituents, alkylsulfonyl, arylsulfonyl
and an aminocarbonylalkyl or aminocarbonyl-
20 alkylcarbonyl group

wherein the aminocarbonylalkyl or
aminocarbonylalkylcarbonyl nitrogen is (i)
unsubstituted, or (ii) is the reacted amine of
an amino acid, or (iii) substituted with one or
25 two radicals selected from the group consisting
of an alkyl, hydroxyalkyl, hydroxyheteroaralkyl,
cycloalkyl, aralkyl, trifluoromethylalkyl,
heterocycloalkyl, benzofused heterocycloalkyl,
benzofused heterocycloalkyl, benzofused
30 cycloalkyl, and an N,N-dialkylsubstituted-
alkylaminoalkyl group, or (iv) that carboxamido
nitrogen and two substituents bonded thereto
together form a 5- to 8-membered heterocyclo,
heteroaryl or benzofused heterocycloalkyl ring

that is itself unsubstituted or substituted with one or two radicals independently selected from the group consisting of an alkyl, alkoxy carbonyl, nitro, heterocycloalkyl, hydroxy, hydroxy carbonyl, aryl, aralkyl, heteroaralkyl and an amino group, or

R^6 and R^7 together with the depicted nitrogen atom form a saturated or unsaturated cyclic imide substituent in which the ring contains five to eight atoms; and

R^{20} is (a) $-O-R^{21}$, where R^{21} is selected from the group consisting of a hydrido, C_1 - C_6 -alkyl, aryl, ar- C_1 - C_6 -alkyl group and a pharmaceutically acceptable cation, (b) $-NR^{13}-O-R^{22}$ wherein R^{22} is a selectively removable protecting group and R^{13} is a hydrido, C_1 - C_6 -alkyl or benzyl group, or (c) $-NR^{13}-O-R^{14}$, wherein R^{14} is a hydrido, a pharmaceutically acceptable cation or $C(W)R^{15}$ group, where W is O or S and R^{15} is selected from the group consisting of an C_1 - C_6 -alkyl, aryl, C_1 - C_6 -alkoxy, heteroaryl- C_1 - C_6 -alkyl, C_3 - C_8 -cycloalkyl- C_1 - C_6 -alkyl, aryloxy, ar- C_1 - C_6 -alkoxy, ar- C_1 - C_6 -alkyl, heteroaryl and amino C_1 - C_6 -alkyl group wherein the aminoalkyl nitrogen is (i) unsubstituted or (ii) substituted with one or two substituents independently selected from the group consisting of an C_1 - C_6 -alkyl, aryl, ar- C_1 - C_6 -alkyl, C_3 - C_8 -cycloalkyl- C_1 - C_6 -alkyl, ar- C_1 - C_6 -alkoxy carbonyl, C_1 - C_6 -alkoxy carbonyl, and C_1 - C_6 -alkanoyl radical, or (iii) wherein the amino C_1 - C_6 -alkyl nitrogen and two substituents attached thereto

form a 5- to 8-membered heterocyclo or heteroaryl ring.


101. The compound according to claim 91
5 wherein D is -F or -NO₂.

102. The compound according to claim 91
wherein R²² is a 2-tetrahydropyranyl group.

103. The compound according to claim 91
10 wherein R², R⁶, and R¹³ are hydrido.

104. The compound according to claim 94
wherein R⁷ is R^{7A}, and R^{7A} is selected from the group
15 consisting of an alkylcarbonyl, alkoxycarbonyl,
arylcarbonyl, alkylsulfonyl, arylsulfonyl,
aminoalkylcarbonyl, aminoaralkylcarbonyl,
alkoxycarbonylamino-alkylcarbonyl,
alkoxycarbonylaminoaralkylcarbonyl,
20 cycloalkoxycarbonylaminoalkylcarbonyl,
cycloalkoxycarbonylaminoaralkylcarbonyl,
aryloxycarbonylaminoalkylcarbonyl, aryloxycarbonyl-
aminoaralkylcarbonyl, aralkyloxycarbonylamino-
alkylcarbonyl, aralkyloxycarbonylaminoalkylcarbonyl,
25 heteroaryloxycarbonylaminoalkylcarbonyl and a
heteroaryloxycarbonylaminoaralkylcarbonyl group.

105. A compound that corresponds in
structure to the formula below



wherein

- 5 R^1 is a substituent that contains an optionally substituted 5- or 6-membered carbocyclo, heterocyclo, aryl or heteroaryl radical bonded directly to the depicted SO_2 -group that has a length greater than that of an extended pentyl group and
- 10 less than about that of an extended eicosyl group, wherein R^1 defines a three-dimensional volume upon rotation about an axis drawn through the SO_2 -bonded 1-position and the 4-position of a 6-membered ring radical or drawn through the SO_2 -bonded 1-position
- 15 and the center of 3,4-bond of a 5-membered ring radical, the widest dimension of said volume in a direction transverse to the axis of rotation being about that of one furanyl ring to about that of two phenyl rings;
- 20 R^2 is selected from the group consisting of a hydrido, C_1 - C_4 hydrocarbyl, hydroxy- C_1 - C_4 hydrocarbyl, C_1 - C_4 hydrocarbyloxy, halo- C_1 - C_4 hydrocarbyl, C_1 - C_4 hydrocarbyloxymethyl, aryl, aryl C_1 - C_4 hydrocarbyl, aminomethyl, (N- C_1 - C_3
- 25 hydrocarbyl)aminomethyl, (N,N-di- C_1 - C_3 hydrocarbyl)aminomethyl, (N-morpholino)methyl, (N-pyrrolidino)methyl, and a (N-thiomorpholino)methyl group;

R⁶ and R⁷ are independently selected from the group consisting of a hydrido, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, alkylaminocarbonyl, aralkyl, carboxyalkyl, carboxyalkylcarbonyl, 5 alkoxycarbonylaminoalkylcarbonyl, heterocyclocarbonyl, heterocycloalkylcarbonyl, aryloxycarbonylaminoalkylcarbonyl, aralkyloxycarbonylaminoalkylcarbonyl, perfluoroalkylcarbonyl, trifluoromethylalkylcarbonyl, 10 alkylthioalkylcarbonyl, arylthioalkylcarbonyl, aralkylthioalkylcarbonyl, heteroaralkylthioalkylcarbonyl, or a sulfoxide or sulfone of any of said thio substituents, alkylsulfonyl, arylsulfonyl and an aminocarbonylalkyl or aminocarbonyl- 15 alkylcarbonyl group

wherein the aminocarbonylalkyl or aminocarbonylalkylcarbonyl nitrogen is (i) unsubstituted, or (ii) is the reacted amine of an amino acid, or (iii) substituted with one or 20 two radicals selected from the group consisting of an alkyl, hydroxyalkyl, hydroxyheteroaralkyl, cycloalkyl, aralkyl, trifluoromethylalkyl, heterocycloalkyl, benzofused heterocycloalkyl, benzofused heterocycloalkyl, benzofused 25 cycloalkyl, and an N,N-dialkylsubstituted-alkylaminoalkyl group, or (iv) that carboxamido nitrogen and two substituents bonded thereto together form a 5- to 8-membered heterocyclo, heteroaryl or benzofused heterocycloalkyl ring 30 that is itself unsubstituted or substituted with one or two radicals independently selected from the group consisting of an alkyl, alkoxycarbonyl, nitro, heterocycloalkyl,

hydroxy, hydroxycarbonyl, aryl, aralkyl, heteroaralkyl and an amino group, or

R⁶ and R⁷ together with the depicted nitrogen atom form a saturated or unsaturated cyclic imide substituent in which the ring contains five to eight atoms; and


R²⁰ is (a) -O-R²¹, where R²¹ is selected from the group consisting of a hydrido, C₁-C₆-alkyl, aryl, ar-C₁-C₆-alkyl group and a pharmaceutically acceptable cation, (b) -NR¹³-O-R²² wherein R²² is a selectively removable protecting group and R¹³ is a hydrido, C₁-C₆-alkyl or benzyl group, or (c) -NR¹³-O-R¹⁴, wherein R¹⁴ is a hydrido, a pharmaceutically acceptable cation or C(W)R¹⁵ group, where W is O or S and R¹⁵ is selected from the group consisting of an C₁-C₆-alkyl, aryl, C₁-C₆-alkoxy, heteroaryl-C₁-C₆-alkyl, C₃-C₈-cycloalkyl-C₁-C₆-alkyl, aryloxy, ar-C₁-C₆-alkoxy, ar-C₁-C₆-alkyl, heteroaryl and amino C₁-C₆-alkyl group wherein the aminoalkyl nitrogen is (i) unsubstituted or (ii) substituted with one or two substituents independently selected from the group consisting of an C₁-C₆-alkyl, aryl, ar-C₁-C₆-alkyl, C₃-C₈-cycloalkyl-C₁-C₆-alkyl, ar-C₁-C₆-alkoxycarbonyl, C₁-C₆-alkoxycarbonyl, and C₁-C₆-alkanoyl radical, or (iii) wherein the amino C₁-C₆-alkyl nitrogen and two substituents attached thereto form a 5- to 8-membered heterocyclo or heteroaryl ring.

106. The compound according to claim 96 wherein R², R⁶, and R¹³ are hydrido.

SUBSTITUTE SHEET (RULE 26)


107. The compound according to claim 97 wherein R^7 is R^{7A} , and R^{7A} is selected from the group consisting of an alkylcarbonyl, alkoxycarbonyl, arylcarbonyl, alkylsulfonyl, arylsulfonyl, aminoalkylcarbonyl, aminoaralkylcarbonyl, alkoxycarbonylamino-alkylcarbonyl, alkoxycarbonylaminoaralkylcarbonyl, cycloalkoxycarbonylaminoalkylcarbonyl, cycloalkoxycarbonylaminoaralkylcarbonyl, aryloxycarbonylaminoalkylcarbonyl, aryloxycarbonylaminoaralkylcarbonyl, aralkyloxycarbonylaminoalkylcarbonyl, aralkyloxycarbonylaminoaralkylcarbonyl and a heteroaryloxycarbonylaminoalkylcarbonyl and a heteroaryloxycarbonylaminoaralkylcarbonyl group.

108. The compound according to claim 98 wherein said compound corresponds in structure to the formula below




20


109. The compound according to claim 98 wherein said compound corresponds in structure to the formula below




110. The compound according to claim 98 wherein said compound corresponds in structure to the formula below




5 111. The compound according to claim 98 wherein said compound corresponds in structure to the formula below




10 112. The compound according to claim 98 wherein said compound corresponds in structure to the formula below




15 113. The compound according to claim 98 wherein said compound corresponds in structure to the formula below




114. The compound according to claim 98 wherein said compound corresponds in structure to the formula below




5 115. The compound according to claim 98 wherein said compound corresponds in structure to the formula below



10 116. The compound according to claim 98 wherein said compound corresponds in structure to the formula below



15 117. The compound according to claim 98 wherein said compound corresponds in structure to the formula below



INTERNATIONAL SEARCH REPORT

national Application No

PCT/US 00/09807

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C317/48 C07C317/50 C07C323/65 C07D211/62 C07D213/56
C07D213/81 C07D213/82 C07D215/50 C07D295/15 C07D307/68
C07D317/58 C07K5/062 A61K31/10 A61P7/00 A61P11/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C C07D C07K A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

BEILSTEIN Data, WPI Data, EPO-Internal, PAJ, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 780 386 A (HOFFMANN LA ROCHE) 25 June 1997 (1997-06-25) page 3 -page 4; examples 24-30 page 76, line 58 --- -/--	1,3-6, 16,17, 32,33, 35-37, 45,94-97

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

26 September 2000

Date of mailing of the international search report

13/10/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

English, R

INTERNATIONAL SEARCH REPORT

national Application No
PCT/US 00/09807

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A61P29/00 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	EP 0 994 104 A (ONO PHARMACEUTICAL) 19 April 2000 (2000-04-19)	1, 3-6, 25, 32, 33, 35, 36, 80, 81, 90, 94-97
X	page 3; examples 12-20 claim 8, compounds 18-20, 22, 23, 25, 26, 28-30, 32, 33, 35-39 & WO 97 49679 A 31 December 1997 (1997-12-31)	
X	WO 98 03164 A (MONSANTO) 29 January 1998 (1998-01-29) page 169	1, 3-6, 32-34, 36, 80

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

Date of mailing of the international search report

26 September 2000

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

English, R

INTERNATIONAL SEARCH REPORT

National Application No

PCT/US 00/09807

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
X	G. KOENIG: "Die Oxydationsproducte der Mercaptursäuren" HOPPE-SEYLER'S ZEITSCHRIFT FÜR PHYSIOLOGISCHE CHEMIE, vol. 16, 1892, pages 525-551, XP000912177 Walter de Gruyter, Berlin, DE ISSN: 0018-4888 page 533 -page 534 ---	1,3-6, 32,33, 35,36, 38,80
X	US 5 480 887 A (W.J. HORNBACH, ET AL.) 2 January 1996 (1996-01-02) column 34, line 64 column 36, line 12 - line 13 ---	100,101, 103,104
X	L. GOODMAN, ET AL.: "Potential anticancer agents. V. Some sulphur-substituted derivatives of cysteine" JOURNAL OF ORGANIC CHEMISTRY, vol. 23, no. 9, 15 October 1958 (1958-10-15), pages 1251-1257, XP002148407 American Chemical Society, Washington, DC, US ISSN: 0022-3263 compound XXXVIII ---	100,101, 103,104
X	R.K. DUA, ET AL.: "S-Aryl-L-cysteine S,S-dioxides: design, synthesis, and evaluation of a new class of inhibitors of kynureninase" JOURNAL OF THE AMERICAN CHEMICAL SOCIETY, vol. 115, no. 4, 24 February 1993 (1993-02-24), pages 1264-1270, XP002148408 American Chemical Society, Washington, DC, US ISSN: 0002-7863 page 1269, right-hand column, paragraph H -----	100,101, 103,104

FURTHER INFORMATION CONTINUED FROM PCT/SA/ 210

Continuation of Box I.2

Claims Nos.: 1,3-6,31-33,35-36,38-44,80,94-97,99,100,102-107 (partially searched), 2,34,98 (not searched)

Present claims 1-6,31-36,38-44,80,94-100,102-107 relate to compounds and their therapeutic effects (see note in Box I, section 1) by reference to the following parameters:

P1: the length and three-dimensional volume of the R1 substituent (claims 1,33,94)

P2: the ability of the substituent D to act as a nucleophilically-displaceable leaving group (claim 100)

The use of these parameters in the present context is considered to lead to a lack of clarity within the meaning of Article 6 PCT. It is impossible to compare the parameters the applicant has chosen to employ with what is set out in the prior art. The lack of clarity is such as to render a meaningful complete search impossible.

Consequently, the search has been restricted to the compounds in the examples and the compounds in claims 7-30,46-58,60-76,78,79,81-93,108-117. Thus the search with respect to compounds was limited to compounds or formulae A, B1 and D in which R1 is a phenyl group substituted in the 4-position by a phenoxy, phenylthio, methoxy or benzoylamino group or a bromine atom or in which R1 is a 3,4-dimethoxyphenyl group and to compounds of formula C in which D is a nitro group or a fluorine atom. Similarly the search with respect to therapeutic effects (claims 94-97,99) was limited to the effects of compounds of formula B in which R1 is a phenyl group substituted in the 4-position by a phenoxy, phenylthio, methoxy or benzoylamino group or a bromine atom or in which R1 is a 3,4-dimethoxyphenyl group.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

information on patent family members

National Application No

PCT/US 00/09807

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0780386 A	25-06-1997	AU 700725 B	14-01-1999
		AU 7548296 A	31-07-1997
		BR 9606134 A	03-11-1998
		CA 2193178 A	21-06-1997
		CN 1160045 A	24-09-1997
		CZ 9603740 A	14-01-1998
		HR 960612 A	28-02-1998
		HU 9603494 A	30-11-1998
		JP 2921673 B	19-07-1999
		JP 9249638 A	22-09-1997
		NO 965413 A	23-06-1997
		NZ 299941 A	27-05-1998
		PL 317604 A	23-06-1997
		TR 970547 A	21-07-1997
		US 5932595 A	03-08-1999
EP 0994104 A	19-04-2000	WO 9749679 A	31-12-1997
WO 9803164 A	29-01-1998	AU 714687 B	06-01-2000
		AU 3890497 A	10-02-1998
		BR 9710758 A	17-08-1999
		CN 1230884 A	06-10-1999
		CZ 9900167 A	16-06-1999
		EP 0939628 A	08-09-1999
		NO 990262 A	17-03-1999
		PL 331369 A	05-07-1999
		US 6013649 A	11-01-2000
US 5480887 A	02-01-1996	US 5624934 A	29-04-1997
		AU 700417 B	07-01-1999
		AU 7930294 A	21-08-1995
		CA 2182192 A	10-08-1995
		CN 1145587 A	19-03-1997
		EP 0744948 A	04-12-1996
		JP 9509155 T	16-09-1997
		WO 9520962 A	10-08-1995

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
30 November 2000 (30.11.2000)

PCT

(10) International Publication Number
WO 00/71514 A1

(51) International Patent Classification⁷: C07C 317/48,
317/50, 323/65, C07D 211/62, 213/56, 213/81, 213/82,
215/50, 295/15, 307/68, 317/58, C07K 5/062, A61K
31/10, A61P 7/00, 11/00

(21) International Application Number: PCT/US00/09807

(22) International Filing Date: 17 May 2000 (17.05.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/134,996 20 May 1999 (20.05.1999) US

(71) Applicant (for all designated States except US): G.D.
SEARLE & CO. [US/US]; Corporate Patent Department,
P.O. Box 5110, Chicago, IL 60680 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BARTA, Thomas,
E. [US/US]; 1133 Maple Avenue, 3W, Evanston, IL 60202
(US). BECKER, Daniel, P. [US/US]; 1800 Maplewood
Lane, Glenview, IL 60025 (US). BEDELL, Louis, J.
[US/US]; 1832 E. Camp McDonald Road, Mt. Prospect,
IL 60056 (US). DE CRESCENZO, Gary, A. [US/US];
7345 Spruce Hill Court, St. Charles, MO 63304 (US).
FRESKOS, John, N. [US/US]; 7572 York, Clayton, MO
63105 (US). GETMAN, Daniel, P. [US/US]; 66 Sunny
Hill Court, Chesterfield, MO 63017 (US). HEINTZ,
Robert, M. [US/US]; 603 Nancy Place, Ballwin, MO
63021 (US). HOCKERMAN, Susan, L. [US/US];

5318 W. Hutchinson, Chicago, IL 60641 (US). LI, Hui
[US/US]; 322 Jefferson Court, Vernon Hills, MO 60061
(US). MISCHKE, Brent, V. [US/US]; 25 White River
Lane, Defiance, MO 63341 (US). VILLAMIL, Clara, I.
[US/US]; 813 Long Road, Glenview, IL 60025 (US).

(74) Agents: WILLIAMS, Roger, A. et al.; G.D. Searle & Co.,
Corporate Patent Department, P.O. Box 5110, Chicago, IL
60680-5110 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE,
DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU,
ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,
LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ,
PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT,
TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— With international search report.

(48) Date of publication of this corrected version:


22 February 2001

(15) Information about Correction:

see PCT Gazette No. 08/2001 of 22 February 2001, Section
II

[Continued on next page]

(54) Title: α -AMINO- β -SULFONYL HYDROXAMIC ACID COMPOUNDS



(57) Abstract: A family of molecules of formula (A) in which R^1 is a 5- or 6-membered carbocyclo, heterocyclo, aryl or heteroaryl radical such that its size defined by certain parameters; R^{20} is -O- R^{21} , -NR¹³-O- R^{22} or -NR¹³-O- R^{14} ; and the other variables are as defined in the claims, is disclosed that inhibit matrix metalloproteinase (MMP) activity, and particularly inhibit the activity of one or more of MMP-2, MMP-9, or MMP-13, while generally exhibiting little activity against MMP-1. A contemplated compound also exhibits little inhibition of the production of TNF. Also disclosed are process for preparing a contemplated compound and for treating a mammal having a condition associated with pathological matrix metalloproteinase activity.



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.